Rebound:

JAN 16 1941

LIBRARY
OF THE
School of Medicine



University of Maryland

Md.
Coll.

UNIVERSITY OF MARYLAND

THESES

1828

| | | |
|---|---|---|
| Merriken, Lachariah | Apoplexy | 22p. |
| Hudnall, Alfred | Inaugaral Dissertation | 27p. |
| Handy, Littleton N. | Cantharides | 45p. |
| Grimes, W, H. | An Inquiry into the Nature, Cause and Cure of Mental Derangement | 33p. |
| Ward, James R. | Yellow Fever | 25p. |
| Grieves, Horatio G. | Phthisis Pulmonalis | 59p. |
| Wharton, John O. | Cynanche Trachealis | 31p. |
| Bussey, Bennett | Pulmonary Consumption | 17p. |
| Hughes, Charles F. | Hydrocephalus Internus | 39p. |
| Lowndes, Leonard | Hydrocephalus Internus | 17p. |
| Goldsborough, Leander W. | Pneumonia | 35p. |
| Clendenin, W. H. | Atmosphere | 65p. |
| Berry, John | Hydrocele of the Tunica Vaginalis Testis | 21p. |
| Stuart, Joseph N. | Tetanus | 35p. |
| McCeney, Edward | Icterus | 29p. |
| Wilson, Paca | Causes of Sudden Death and the Means of Preventing It | 30p. |
| Kirk, Wm. N. | Inflammation | 21p. |
| Carmichael, George F. | Necrosis | 23p. |

R45

1

# Apoplexy

An Inaugural Dissertation

by

Zachariah Merriken

Annapolis Maryland

Spring 1828

In speaking of Apoplexy, the disease of which I am about to treat, I shall not pretend to claim originality. The disease has been called by several different names, or more properly, several diseases have been confounded with Apoplexy viz^t. Lethargus, Cataphora, Carus, and Coma, these are supposed to be the same disease, differing in degree only; but of these it is not my intention to speak, therefore, I shall confine myself strictly to Apoplexy. The term Apoplexy has been given the disease from the suddenness of its attack, it is derived from the Greek words απο which signifies upon and πλησσω to strike suddenly. In this disease the whole of the external and internal senses and the whole of voluntary motion, are in great degree abolished, while respiration and

the action of the heart continues to be performed it has also been called by the Latins Morbus Attonitus they have also called it Lideratio from a belief that it was produced by planatary influence

Dr. Cullen has placed this genus of disease in the class Neuroses and the order Comata he has also taken notice of nine Genera of this disease besides those which are symptomatic. First Apoplexia Sanguinea and this he defines to be universal Plethora with a determination of blood to the head Second Apoplexia Serosa which generally happens in old and Leucophlegmatic persons. Third Apoplexia Hydrocephalica in this genus the disease is produced by the pressure of water upon the brain Fourth

Apoplexia Atrabilaria observed in Atrabilarious or melancholic temperaments Fifth Apoplexia Traumatica when the head is injured by violent external force Sixth Apoplexia Venenata from poisonous matters whether taken internally or applied externally Seventh Apoplexia Mentalis from passions of the mind Eighth Apoplexia Cataleptica in which respiration is not stertorous and though the limbs maintain any occasional position give way to force applied to them Ninth Apoplexia Suffocata as happens from hanging or drowning. The division of the disease into so many genera tends more to confuse than to instruct the young practitioner. Apoplexy most frequently attacks persons at an advanced age and Dr. Cullen says especially

those above sixty years of age but a great majority of the cases which I have seen have occurred in persons under fifty and in those of short necks and plethoric habits and who have passed a sedentary life and used a full diet Those who have been intemperate both in ~~habit~~ diet and drink or those who have indulged in one and not in the other have been attacked by the disease The sudden suppression of an accustomed Haemorrhage The most frequent precursory symptoms are fits of giddiness, head ache hæmorrhage from the nose some transitory interruption of vision and hearing and these two senses are also false in some cases a faultering of the tongue frequent drowsiness and repeated attacks of incubus but these precursory symptoms are by no means uniform ~~but~~

for the disease in some cases suddenly attacks the patient without these symptoms having appeared but when those symptoms do occur they enable us to foresee the impending danger and to avert its fury by a proper attention to the present state of the patients system and by a due regard to regimen we may enable and if not entirely to escape this too fatal disease at least to render its attack longer than he otherwise would have done. When the disease has come on suddenly it has been frequently observed to have been induced by violent exercise this remark I have seen verified in the case of a Farmer who though for some time previous to his attack had led a very active life: had in the earlier part of his life been quite sedentary in consequence of his avocations at that time and had in the whole

of his life been a very free liver and by these two latter causes had acquired a predisposition to the disease he was of a plethoric habit answering very well the description already given of those most liable to the disease This gentleman after a severe exercise of almost a whole day was suddenly attacked in the absence of any premonitory symptoms of the approach of this malady. Although says Dr. Cullen the whole of the body is affected with the loss of sense and motion it sometimes takes place more upon one side than on the other and in this case the side least affected is frequently convulsed. Stertorous breathing is often present and has been supposed to indicate the violence of the disease but this is denied as it has not at all been present in a complete and most violent

state of the disease. After having thus enumerated most of the principle symptoms of the disease we will now advert to the causes and first to the proximate cause which has been defined to be whatever interrupts the motion of the nervous power of the brain to the muscles thereby destroying voluntary motion or so far as sense is affected whatever interrupts ~~the sentient extremities~~ the motion of the nervous power from the sentient extremities of the nerves to the brain. Such interruptions are said to be produced by some compression of the origin of the nerves or by something destroying the mobility of the nervous power the loss of sense and motion in particular parts of the body may be occasioned by a compressio

of the origin of the nerves going to that part or by compression of the same nerves at a point intermediate between their origin and the part diseased by such compression. Compression of the origin of the nerves may be produced in different ways viz: by a depression of a part of the Calvarium. A congestion of the vessels of the brain has also been said to have produced it but on this subject there is a contrariety of opinion. It is Dr. Nathaniel Potter's opinion that this is a frequent cause of the disease. It has on the contrary been maintained by Dr. Davidge as well as other distinguished Physiologists that a congestion of the vessels of the brain never does take place. It may be produced by tumours either of the bone or soft tissue.

If compression of the origin of the nerves be from a depressed portion of the brain case it is to be removed but this operation more properly appertains to surgery The consideration of bony tumours and also of fleshy tumours may be omitted as in most instances they are neither to be discerned nor cured by any means. The effused fluids are of two kinds they may be either a portion of the common mass of the blood poured out by vessels carrying red blood or it may be serum or a fluid resembling serum which is poured out by the exhalants. This fluid alluded to resembling serum has been supposed not to be serum but to be a fluid changed by the action of the small vessels from which it escapes however be it serum or a changed fluid it makes no change of treatment in the disease these effusions have been said to be para-

duced by whatever ~~can~~ increases the impetus of the flow of blood in the arteries of the head the same cause has been assigned to the congestion of the blood vessels of the brain this is a probable cause if congestion of them does ever take place but as there is a difference of opinion on this subject among the most enlightened Physiologists it therefore cannot be expected that one of my inexperience would presume to hazard an opinion in opposition to either of the parties who thus dissent. An impediment in the free return of blood from the vessels of the brain has also been enumerated among the causes of effusion of serum. There are two other causes of this serous effusion. The one is a relaxation of the exhalants. The second is an over proportion of watery parts in the blood.

which is therefore ready to rush of by the exhalants but whether compression be produced by a serous or a sanguine effusion the indication of cure are the same or nearly so. so besides these, other causes are taken notice of by authors as the Mephitical arising from fermenting liquors the fumes which arise from mercury lead and other metalic substances have produced it. The modus opperandi of those noscious poisons is I think very correctly given in the 111th Lec: of the 1st lines of the practice by Dr Cullen and I think it will not be amiss here to transcribe it I believe says the Dr their immediate and direct action to be on the nervous power destroying its mobility because the same poisons shew their power in destroying the

irritability of the muscles and of the nerves connected with them when both of these are entirely seperate from the rest of the body. Having thus taken notice of the chief causes of Apoplexy and of its symptoms I shall now proceed to give the treatment when there is evidently a predispositions to the disease all exciting causes ought carefully to be avoided the most common causes of the disease have been already stated the immoderate use of spirituous liquors ought also to be carefully avoided an indulgence of the appetite in high seasoned dishes must be restricted in short all the causes calculated to induce plethora should be avoided but when this Plethora or in other words this predisposition to Apoplexy is either hereditary or is acquired by intemperance in eating or drinking remedies

should be employed for its cure if it be produced by intemperance the cause should be removed and blood letting and abstinence from animal food should be enforced. The mode of bloodletting has been considered of importance some have recommended it to be taken from the Jugular vein from the temporal Artery &c. The apperation of Venæ Section has also been thought more effectual when performed in the side opposite to that most affected but I think for the sake of dispatch the arm most convenient to the Physician should be bled from. Purging is also a very important remedy this should be attempted by stimulating injections and if the power of deglutition remain active Cathartics should be administered Emetics have been recommended but these have been

objected to on the presumption that it would impel the blood to the brain with so much violence as to rupture the vessels or if they are already ruptured to cause a more copious effusion where the stomach contains a hearty meal it will generally be ejected from the stomach. if a considerable quantity of blood be drawn from the arm or jugular vein or from any convenient and safe vessel and in this case very beneficial effects have resulted. blistering has been recommended but I think an important remedy in the early stage of the disease stimulants have been almost universally employed but I think these might be employed advantageously

in cases of suspended animation but do not think them useful or their employment warrantable in Apoplexy as the violent efforts in vomiting will as before stated impel the blood to the brain with so much violence as to increase the effusion or rupture the vessels which are not yet ruptured. If a poison introduced into the stomach produces Apoplexy and if a spontaneous vomiting occurs it should be encouraged by draughts of tepid water and by the remedies usually recommended in cases of poison. If stimulants in the later periods of the disease become necessary they are to be employed and the most proper will naturally be-

resorted to by the skillfull and prudicious practitioner The affusion of cold water has been thought to be of much value in rousing a apoplectic from the insensible state. When there is a full habit or a plethora existing the pen if applied to the back of the neck has been useful in overcoming this predisposition so also is a seton introduced into the back of the neck these are merely used for the prevention of Apoplexy but when Apoplexy seriously threatens there is nothing which can be substituted for copious Venæ section. If Apoplexy actually does occur bleeding is immediately to be employed and

if as above stated the powers of deglutition remain
to the patient cathartics should be administered
and their apperation promoted by injections.
cuping the temples also leeching may be emp
loyed with good effect. rest should be strenuously
enforced the head being considerably raised by
pillows above the level of the body and if the
patient recovers a total avoidance of the pre
disponent causes and likewise of plethora should be
strictly adhered to by the patient. A paroxysm
of deep intoxication has been confounded with
Apoplexy but may be distinguished ~~by the paroxysm~~
generally by the history of the case and by the
smell of liquor being perceptible in the

breath deep sleep resembles it in some degree but is readily distinguished by the patient being aroused by the application of volatiles applied to the nose by pinching or by calling him in a loud tone of voice it is also confound with Palsy and Syncope but is distinguished from the former by its being an affection of the whole of the powers of sense and voluntary motion and from Syncope by its being attended with the continuance of respiration and the action of the heart,

An Inaugural Dissertation
Submitted to the examination
of the Provost and medical
Professors of the University of
Maryland,
by their friend
Alfred Sudnall
of Northumberland County Virginia
in the year of our Lord A.D. 5828. A.D. 1828.

24

The following lines are respectfully dedicated to the Medical Professors in the University of Maryland — as a small tribute of regard and esteem entertained for them, both as private individuals and public teachers

by their friend —

Alfred Hildreth

26

## Remarks on Tetanus.

The disease which I am about to give a discription of, is one which has in former times excited terror in every medical Practitioner, who has the misfortune to witness a case of it.—
I mean Tetanus— A genus of disease, in the Class Neuroses, and order Spasmi, of Cullen, characterised by a spasmodic rigidity of almost the whole body.——— The varieties of this Disease which are discribed by Dr Cullen are, First, Opisthotonos when the body is violently drawn backwards, by the contraction of the Dorsal muscles— Secondly, Emprosthotonos, when the body is bent forwards— Thirdly, Pleurosthotonos, when the body is drawn to either side and Fourthly, Trismus, or locked Jaw.——— Though I consider them one, and the same disease and these distinctions of course unnecessary.

## Causes.

Tetanic complaints may arise from many and various causes such as, punctures, lacerations, amputations and fractures of the Limbs— Gunshot wounds, wounds in the palms of the hands and soles of the feet by means of nails, or splinters of wood,— the Injury done to the feet by Frost, cutting the umbilical cord in new born Infants, and the sudden application of cold to the body after it has been exposed to extreme heat— It has also been sometimes induced by certain

28

poisonous vegetables received into the stomach — as Hemlock — Stramonium &c. ——— This Disease is said to occur most frequently in warm climates, and in the warmest weather of those climates, but this is by no means a general rule, for we sometimes have it in our winters.

This Disease affects all ages, sexes, Temperaments and Complexions — but the nervous and irritible are most subject to it. ——— Some have said that negroes are more predisposed to attacks of it, than white people they are more frequently affected with it but this, probably arises from their being more exposed to injuries than the whites, and not from predisposition, they are more liable to tetanus, because they are more subject to injuries as they generally go barefooted and from that circumstance are very often wounded about the feet with nails, splinters of wood, broken glass &c. &c. —

## Symptoms. —

This Disease sometimes comes on suddenly to a very violent degree but more generally it approaches by slow degrees to its violent stage.

It commences with a sense of stiffnes in the back part of the neck which, goes on increasing until it renders the motion of the head difficult and painful — After the rigidity of the neck has come on there is commonly at the same time a sense of

uneasiness felt about the root of the tongue - which by degrees generally becomes a difficulty of swallowing and if, the disease is not combated properly at length an entire interruption of it.
While the rigidity of the neck goes on increasing there arises a pain often very violent at the scrobiculus Cordis, and shooting from thence into the back —— When this pain arises all the muscles of the neck and particularly those of the back part of it are affected with spasm pulling the head strongly backward At the same time the masticating muscles are affected and the jaws are violently clenched, so that it is sometimes necessary to extract a tooth to introduce any thing into the stomach. —— This is what writers have named Locked jaw and is very troublesome; the tongue is sometimes very much injured by the teeth in consequence of its being thrust out of the mouth during the interval of spasm and it is in that way that it is so often injured. —— When the disease has advanced thus far the pain at the bottom of the sternum returns very frequently — and with it the spasm of the back part of the neck. As the disease thus proceeds a greater number of muscles come to be affected

with spasm.— In succession to those of the neck, and jaws, those of the back bending the whole body strongly backwards and this, is what has been called Opisthotonos.—
The muscles of the superior extremities are not the only muscles affected, for both the flexor and extensor muscles of the legs are very rigid, and the limbs are very stiff.—
During the whole of the disease the abdominal muscles are violently affected with spasm, so that the abdomen is strongly retracted and feels very hard.
At the height of the disease every organ of voluntary motion seems to be affected; and among the rest the muscles of the face: The forehead is drawn up in furrows, the eyes are sometimes distorted, are commonly rigid and immoveable in their sockets; The cheeks are drawn backwards towards the ears so that the whole countenance expresses the most violent grining, under these universal spasms a violent convulsion commonly comes on and puts an end to life.
The attacks of this Disease are seldom attended with high fever although there may be occasionally some:— And it is said, by some that at the commencement of this Disease the pulse is

often full and frequent, tense and strong.—
But I believe in the majority of cases,
there is no fever of consequence; and most
commonly the pulse is weak and quick.
In this Disease the head is seldom affected
with delirium, or confusion of thought till the
last stage of it.
In this disease I believe the natural functions
are not either immediately, or considerably
affected, however occasionally there is vomiting
in the early part of the disease but, this
generally continues a very short time;
and it is quite usual for the appetite
of hunger to remain through the whole course
of the disease, and the food that is taken,
seems to be regularly digested, however this
is in very small quantity.
The excretions are sometimes affected but not
always— The urine is sometimes suppressed, or is
voided with difficulty.
The belly is sometimes costive this I think
originates from opiates, for we have no accounts
of it unless opiates were used.
This disease until very lately generally proved
fatal, this was owing very much to the manner
in which patients were treated by the older
practitioners; For we know that until very lately

Physicians were not acquainted with a method of cure and that, since a more proper method has been known and carried into effect, many have recovered:— It may be concluded that the fatal tendency of it is not so unavoidable as has been supposed.—

In judgeing of the disease in particular cases, we may observe, that, if it comes on very rapid it is more severe than when it comes on more slowly.— (The first case of Tetanus, which I saw terminated fatally on the fourth Day.)

This disease if not properly treated, often proves fatal before the fifth, or sixth, Day, and when a patient has passed this period he may be said to be in greater safety, and in general the disease is the safer, the longer it has continued.

It is to be observed however that the Disease is dangerous for many Days after the sixth.

## Treatment.

I come now to speak of the treatment which should be carried into effect with perseverance, and experience has taught us that Opium, and Mercury, has often proved an effectual remedy, but to render them so they must be given in large quantities so that we may have their full effects.

I am fully persuaded that opium is very seldom given in sufficient doses.
The practice has been, to give it either in a liquid, or solid form, not in very large doses at a time, but in moderate doses every one, two, or three hours; this practice will do if its effects are not suffered to go off, however it does not seem to opperate in this Disease, as it does in others; For although it procures some remission of the spasm, and pain, it hardly ever induces any sleep, or occasions that stupor, or delirium which it often does in other circumstances, when much smaller quantities have been given.
It is therefore observed that in this affection there should be no retailing, that it should be given by wholesale and it should be given as largely, and as frequently, as the symtoms of the Disease may seem to demand.
It must however be observed that though the first exhibition of the opium may have produced some remission of the symtoms yet, the effects of it do not long continue in the system and this disease being very ready to return, it is very necessary by the time that the effects of the opium given, are going off and especially upon the

least appearance of the return of the spasm, to repeat the dose in the same quantities as before.— This practice should be very strictly pursued as long as ever the disease showed any disposition to return: And it is not until after the Disease has already subsisted for some time and when considerable and long continued remissions have taken place, that the doses of the Opium may be diminished and the intervals of exhibiting them, be more considerable.

Opium administered in this way, has in many cases been successful, and probably would have been so in many others if, it had not been too sparingly used, either from timidity of Practitioners or from its exhibition being prevented by that interruption of deglution which so often attends this disease. Which latter circumstance requires that, the medicine should be immediately and largely employed upon the very first approach of the disease, before the deglutition becomes difficult or if this opportunity be lost the medicine in sufficient quantity and with due frequency should be thrown into the rectum by glysters.

It has been recommended to employ opium with

some of the antispasmodics, such as musk, and camphor, this I think is a very weak practice and would answer a very little better purpose than opium alone, in one case I see opium and musk combined but it terminated fatally. Blisters to the neck were also used.— What would be the effect of Emetics in this Disease — We know, that they produce powerful effects sometimes in relaxing the muscles in spasmodic and convulsive Disorders.

The warm bath has also been spoken of as a remedy.

Purgatives has been very highly recommended by some of the Continental Practitioners.

I do not know how they would answer as I have never seen them used.

Bloodletting should not be used except in plethoric subjects.

Blistering was at one time very much used, it is asserted by some Practitioners that it is always hurtful.

Moxa has been used with decided advantage in late years, it should be applied all down the spine, and the eschars dressed with mercurial Ointment— so as to keep up a discharge for some time

Cold bathing has also been recommended by the Continental Practitioners.

I am induced to believe that not one half

of the remedies recommended are of any service. I think that wine, Opium, Moxa, and Mercury are the principal remedies to be relied on in this Disease. — Of these remedies the Wine, Opium, and Moxa, retard the progress of the Disease. — Preventing death from exhaustion until the Mercury shall have time to eradicate it, by converting the Tetanic affection into a mercurial action. — However I should strongly recommend Opium and Wine, for without their aid the Mercury would not have time to act, although they are insufficient of themselves without mercury. The Mercury should always have its full effects, and this effect should be kept up for some time, for if it is not the Tetanic affection will certainly return.

The Mercury should be used very extensively, applying it in large quantities in the form of Mercurial Ointment.

Tobacco has also been found useful and spoken very highly off, by Dr Jackson of Northumberland in Pennsylvania — in a case which occured in his Practice — produced by an injury received on the back of the hand by a splinter of bone, the part had healed, and he made an incision into it, and applied a poultice over it, made of a strong decoction of Tobacco and linseed meal. —

He applied it from the hand to the shoulder, and changed it every six hours: a similar application was made to the scrobiculus Cordis – in order, to operate on the whole system.
In six hours, the symtoms had ameliorated, and the same means were continued with the addition of Castor oil to open the bowels – Thirty six hours from the first application, the Tobacco produced some nausia, and so much Vomiting on the third day, that the whole complaint appeared to be subdued, his stomach was now restored by cordiac nourishment – Drink. &c. Huxhams Tinct – About the same time Erysipelas of the arm came on, succeeded by Mania a Potu which rendered the use of opium, Camphor, &c. necessary. – vid. Med. Rec. Oct 1824 page 315

## Prophylactics.

The best preventives of Tetanus consist in the conversion of punctured into incised wounds by means of a scalpel, or some such instrument, and the exciting of suppuration in wounds that, are lacerated.
These precautions should be used immediately after the injury is received, and before the Tetanic symtoms shall have made their appearance if circumstances exist to prevent the dilatation of punctured

wounds the application of some escharotic should be used such as Ley, Corrosive Sublimate, Spirits of Turpentine, Tincture of Cantharides, and Nitrate of Silver.
These should be followed by poultices so as to bring on suppuration.
If the patient cannot swallow, food should be thrown into his stomach by means of a gumelastic tube.
This may be done by removing a tooth or two.

An
Inaugural Essay
On
Cantharides;
Submitted to
The Examination
of the
Provost
The
Trustees & Medical Faculty
of The
University of Maryland
For The Degree of
Doctor of Medicine
By Littleton D. Handy of
Maryland

March 1828.

To

John Buckler, M.D.

Adjunct Professor of Anatomy

In The

University of Maryland

This Production

of

Youthful Thoughts

On

Medicine

Is Dedicated As

A

Tribute of Respect & Esteem

From

His Pupil

The

Author

# Essay on Cantharides

Cantharides when applied to any part of the surface of the body produce inflammation of the skin and in a few hours there is a preternatural discharge from the capillaries. This discharge being accumulated between the cuticle and cutis vera is called blister. A similar evacuation of fluid may be produced by other acrid substances such as sinapism and many others. But no experiments hitherto made have been attended with such success as to entitle them (sinapis) to such general use as the cantharides: and they are seldom had recourse to except when the flies cannot be obtained or where in consequence of the extreme torpor of the system they have been found ineffectual

The ancients were not entirely ignorant of the powerful effects of blisters although their application was confined to very few diseases. It is said that the Arabians were the first to use them with a view to rouse their patients from a lethargic state. The physicians that succeeded them for sometime used them indiscriminately in every state of morbid action in the system. From an ignorance of the proper period in which they were indicated (as it may be supposed) they did great injury by too early an application of them. our knowledge of the efficacy of blisters may be attributed to a dispute which took place between some Italian physicians relative

the to their use in a plague which prev-
-ailed in the sixteenth century. Their
application cannot be considered as
yet regulated in all cases they are
even now often prematurely applied
and seldom without manifest injury
to the patient. In the application of
blisters we have a very convincing illustr-
-ation of the necessity of attending to the
state of the system before we prescribe
a remedy. This principle is inculcated
with much ardour by a learned and
ingenious professor of the University of
Maryland — upon this principle is
founded the art of alleviating and
curing diseases and as such should never
be forgotten by a practitioner of
medicine — Much has been said about-

the manner in which blisters operate one party maintaining that an absorption of the cantharides takes place and that all their good effects are derived in this way — Another that they act by revulsion only. It appears that the former party were led to their opinion by observing that strangury is produced by the absorption of the acrid particles of the cantharides, This fact they conceive to be sufficient That some part of the cantharides is sometimes taken into the system by means of the absorbents is very certainly proved by the phenomenon of strangury — The absorbed part appears to be particularly determined to the urinary organs sometimes producing by its irritating action inflammation on the neck of the

bladder and consequently pains & dysury
Blisters are powerfully stimulant. we
may infer this from the increased force
and frequency of the pulse which is prod-
-uced by them. and from the great deter-
-mination of blood to the part which is
in contact with the flies. occasioning infl-
ammation and effusion. When blisters are
applied to the extremities they give a centrif-
-ugal determination to the disease thereby
saving parts essential to life as in yellow
fever. The operation of blisters is both
stimulant and sedative. They answer the
purpose of metastasis better than any
other medicine in use. Dr. Rush tells
us " that it is from ignorance or inatt-
-ention to the proper period or stage of
fevers in which blisters have been applied

that there have been so many disputes among physicians respecting their efficacy when applied in a state of great arterial action they do harm: when applied after that action has nearly ceased they do little or no good." The period in which blisters are useful is called by him the blistering point. Blisters are applicable in local and general diseases. We must keep in view the principle which I have before said is so strongly inculcated by a learned Professor —

In the malignant state of fever which appears in the plague yellow fever &c blisters may be applied to the extremities. after the morbid action has been in some measure

reduced by plentiful evacuations. The determination which they produce to the surface often prevents congestion and inflammation in parts essential to life. They have been used in the plague with great advantage as the irritation which they occasioned invited the disease to parts not absolutely necessary to the vital functions and thereby prevented the two often fatal consequences of congestion in the viscera Dr. Rush in his treatise on yellow fever says that blisters when applied in the proper period of this disease (yellow fever) did great service. This time was when the fever was so much weakened by evacuations that the

artificial pain which is produced by the stimulus of the blisters destroyed and like a conductor conveyed off the natural pain of the body.

In typhus or the low state of fever, blisters may be applied as stimulants with considerable advantage. ~~and~~ To obtain this to the greatest extent they must be small and repeated, and never suffered to discharge long: as the evacuation would counteract the purpose for which they are intended. In the typhoid or slow nervous ~~state~~ fever Dr. Huxham used them (blisters) with advantage, and from their operation in other states of fever we can entertain no doubt of their application being attended

with efficacy in this.

The intermittent fever is sometimes so obstinate in consequence of marsh miasmata producing so great a degree of action in the system that the bark has been found inefficacious and in many cases extremely injurious. when such cases occur the patient's pulse is found to be moderately tense. By premising blood letting or the abstraction of the excitement from the blood vessels to the surface by blisters to the wrists the bark will always be found effectual —

Blisters are very important remedies in the treatment of the

different varieties of phlegmasical diseases.

In acute pulmonic affections especially blisters are recommended. Physicians disagree as to the proper time for employing them in pneumonia. That blisters may be employed early in this disease is proved without doubt however they will be more beneficial after the reduction of the vehement arterial excitement.

In relation to this point Dr. Thomson makes the following remarks: "It has sometimes ~~occurred~~ struck me very forcibly" says he "that the precipitate application of blisters to the

chest before general or local blood ~~letting~~ letting, is a prejudicial practice: at least I have seen hydrothorax rapidly follow it, from the increase of the general and topical excitement which blisters thus applied had apparently produced. This point is therefore worthy of further investigation in the acute pulmonary inflammation;+ In the treatment of hepatitis blisters are very useful; and in inflammation of the bowels and peritoneum they are indispensable. In all these affections the blisters should

+ Practical illustrations on the scarlet fever. P. 154

be large and applied immediately over the parts affected.

In all those affections of the throat known by the names of cynanche tonsilaris, parotidea, maligna, and trachealis blisters are very useful applied in the proper time.

In rheumatism I believe every practitioner can testify to the efficacy of blisters. Their good effects must be ascribed to their producing a new action on the skin which is more violent than the inflammation seated within: and probably the

evacuation which they occasion contributes in some measure to relieve the inflammation of the ligaments by lessening the quantity of the determined fluids to the part—

In gout by determining the inflammation to external parts blisters often prevent morbid congestion from taking place in the viscera—

In the maniacal state of fever as in all other states of morbid action we must be directed by the action of the

blood repeats as to the proper time for using them. for the application of them when the system is highly excited. or too enfeebled, would not be attended with advantage but on the contrary with very great injury.

In the apoplectic, phrenitic and lethargic states of fever, blisters may be applied to the head or neck with advantage. we must consult the state of the system and bring it to the point proper for their application. by evacuation

In the paralytic state of fever blisters are often serviceable when applied to the limb or part affected

In the hydrocephalic state of fever they should be applied behind the ears, or to the back of the neck. ~~If effusion has taken~~ If used before effusion ~~place~~ has taken place, they generally prevent it by producing a determination of the morbid action to the skin, and an evacuation which seldom fails to remove the phlogistic diathesis. but even when we have reason to suppose that effusion has taken

place in the brain. they should be used, as they produce a discharge which often relieves the brain from the pressure of the effused fluid,

Blisters applied to the thighs in the amenorrhagic state of fever produce an increased determination of the blood to the lower extremities and some effusion which seldom fails in lessening the activity of the pulse and restoring the natural evacuation

The hysterical and hypoch-

-ondrical states are not unfre-
-quently accompanied with such
force of action in the blood vessels
as to constitute the proper time
for the application of blisters.
The wrists are the proper places
in such cases for them.
The artificial action they
produce abstracts the excitement
from the blood vessels and by
diverting the attention from the melan-
-cholly subjects, the train of gloomy
associations is broken off —
In the waking or watching

state of fever. blisters may be applied to the coristt. By abstracting the excitement from the blood vessels of the brain and fixing it in the skin and muscles. they often induce sleep

These are some

infantile diseases in which blisters may be used with success

The hooping cough in some instances may be called a very distressing disease. The morbid action is sometimes so virulent

as to require blood letting and blistering to a considerable extent

To obviate or remove the inflammatory determination to the lungs blisters should be applied to the thorax as being more efficacious than when applied to distant parts.

This disease often continues a long time after the contagion has ceased to act. and that from ~~habit~~ the power of habit alone. In this case they have a powerful influence in destroying the force of habit. by the new disease which

they produce. Strangury produced by blistering has been attended with an immediate removal of cough in this disease.

In colera infantum blisters applied over the stomach, are very effectual in removing the sickness at the stomach and suppressing the diarrhea. Their efficacy is more certain combined with the fever of the air of the country. The fever which is sometimes produced by the irritation of the gums in dentition is often cured by a blister

to the back of the neck, or one behind each ear. after evacuations have been made. The difficulty of getting blood from children and of exhibiting sufficient doses of medicines renders Epispastics. not unfrequently the only remedy to be depended on. Blisters are sometimes very efficacious when applied locally. The application of a blister to the tract of an inflamed nerve is a practice of much value. This treatment was first ~~used~~ introduced by Dr. Physick, a small plaister of simple cerate spread on linen. is to be applied to the orifice. and over this a blister large enough to cover the whole inflamed part

extending three or four inches from the orifice in every direction.

In concluding this essay permit me gentlemen. to acknowledge the sentiments of gratitude I feel for the civil and polite attentions received from you individually and the information derived from you collectively as Professors of this institution. I fondly anticipate the day when this University will hold a rank distinguished in the records of medical history where the true principles and practice of the different departments of the science of physic are ably taught and the dogmas of ignorance and prejudice consigned to oblivion

# An Inquiry Into The Nature, Cause, & Cure of Mental Derangement.

Submitted to examination before the Provost, and Professors of the University of Maryland, for the degree of Doctor of Medicine,

by

W. H. Grimes, of Maryland.

1828.

To
Jno Reynolds, M.D.

Sir,

under your
direction I received the first rudiments of med-
icine, & to you I dedicate, ~~the~~ my first effort in the
science.

I am induced to do this by two consider-
ations - first, because it is the only way I have at
present of acknowledging the many obligations I am
under to you: second, because I believe, that you
will view favorably any imperfections the attempt
may labour under.

I remain
Sir
Yours very much
obliged
W. H. Grimes

Maryland Hospital
21st of March 1828

# An Inquiry &c.

In selecting mania as the subject of an inaugural thesis, my object was, not so much to advance a new doctrine, as to establish an old one.

It is not without a considerable degree of "fear & trembling," that I enter upon so difficult & so obstruse a subject; but as my object is truth, I shall persue it, at least, with firmness, if not correctness.

I define mania to be a disease of part, or of all the faculties, of the mind. But before entering upon the inquiry, it may, perhaps, be proper to state what those faculties are; not that I intend to be so presumptive as to attempt to say what is the nature of the mind, but that I may endeavour to ascertain the seat of some of its diseases.

The faculties of the mind, according to some writers, are "understanding, memory, imagination, passions, principle of faith, will, the moral faculty, conscience, & sense of the Deity." Its operations sensation, perception, judgement, volition &c.

The mind is put into operation, by impressions made upon it, thro' the medium of the external senses, as— the different colors make impressions on the organs of vision, odours on the organs of smell, different surfaces on the sense of feeling, sounds on the auditory nerve, sweets & acids on

he organs of taste; thus a communication is preserved between man & the material world. When any of these organs are diseased, false impressions are conveyed to the mind — the mind directs the will improperly, we have incoherant actions: thus, a man imagines he sees a spectre, & he flies to avoid it. A disease in any of the faculties is the first step to madness.

In conducting our inquiry we should be led, as well in this as every other disease, first, to ascertain, if possible, the seat of the disease, second, its pathology, third its causes, & fourth its removal or palliation.

Concerning the seat or proximate cause of mental derangement, there have existed a variety of opinions. The ancients supposed that madness was derived from morbid functions of the liver; & as this viscus is very liable to disease, the opinion received considerable countenance from its being very often found in a morbid condition, on dissection after death. Subsequent writers refer it to the spleen; while more modern ones contend that it cannot be referred to either the one or the other, & that it has its seat in the blood-vessels of the brain, or in excessive arterial action, produced from the same causes, which produce fever. Without stoping to inquire into these different opinions, to ascertain their correctness, or to enter into any metaphysical speculations concerning them; I will merely remark, that, if the disease could, with propriety, be referred either to the liver, or spleen, or bloodvessels of the head, we might be led to expect, from the remedies successfully used in these diseases, more fre-

ent cures of madness, than we are in the habit of wit-
essing.

In rejecting all the theories which have been adduced
this subject, we should only be actuated by the hope of
oducing something which approximates nearer to the
uth; whether we shall succeed or not, rests with future
nvestigators to determine — We shall at least make
step towards it, if we fail in our attempt, we may
erhaps, kindle a spark, by which some more success-
ul genius may fire his torch, & thus penetrate the gloom
hich envelops this intricate subject.

We conceive madness to be entirely, a nervous
disease.

To illustrate this position, we shall form
the nervous system into three divisions viz: —
Those of sensation, in which we believe the disease
to be exclusively situated, when these nerves are in
state of disease, & we can easily imagine an hun-
red causes, which may produce a derangement in
their functions; those of motion, & those of sympathy.

That the disease is thus situated, we infer
I. From a total absence of some of the senses, & a per-
version of the rest in every madman: Thus, one
will tear his blanket & roll it up, then cry most
hiddiously until the "Sea Serpent" is removed from
his cell — Here is evidently a diseased nerve, trans-
mitting false impressions to the brain. Another
is insensible to taste & smell, & will swallow any-
thing presented, even of the most disgusting char-

ter, without the least uneasiness being produced. A
ind is insensible to the temperatures of heat & cold
so remarkable a degree, that he will burn his legs
to blisters," without evincing the least concern at it;
. Rush relates the case of a man in the Pennsylvania
ospital, who would stand in one position, in the cold,
ntil mortification was produced, without manifest-
g the least pain, or disposition to move.

II. From sudden impressions made upon the nerves,
requently removing the disease; such as fear, terror
c. For examples of this nature, consult Rush on the
ind, Crichton on mental derangement, and every
ter writer, who wrote on the subject.

III. From the disease disappearing on the accession of another
hich had a tendency to abstract morbid irritability
rom the diseased nerves: thus, the patient, who perceived
the sea serpent in his blanket, took the confluent
small pox in the Maryland hospital, & died; but pre-
ious to his death he became perfectly sensible, &
spoke as rational as any one.

And VI. From its seldom appearing in persons, who have
not attained the years of maturity. The manner of account-
ing for this, is, first external objects do not pro-
duce the same impressions in youth, that they do in
adult age; 2d impressions made in youth are of a
transient nature, & soon forgotten, whereas in adult age,
the impression, ~~the~~ of a beautiful object is continued un-
til the nerve, on which it is made, becomes diseased, from

ong action: just as the continued impressions from odours, on the olfactories, produce hysteria & other diseases, bordering on madness; or, just as the whole system becomes debilitated from long continued exertion, & then inducing disease by keeping up the exertion.

Causes,

We come now to consider the most frequent causes of mental derangement. These are, hereditary predisposition;— this disposition exists in a latent state, & is transmitted from the parent in the same manner that Gout is; but whether transmitted by bloodvessels, nerves, bones, or muscles, "is a mystery locked up in the strong box of nature;" but that the disease is inherited by the offspring from the parents, is a fact, which, I believe, is not at all questioned now. Injuries of the head, so as to make pressure on the nerves of sensation, a constant habit of intoxication, onanism, extreem absence from action, superstitious dread, religeous fanaticism, all the passions, as, Excessive grief or disappointed love, A peculiarity in the formation of the cranium. out of all the maniacal patients now confined in the Maryland Hospital, more than two thirds of them have some peculiar formation of the skull. Some have remarkably small round heads, while others have long narrow heads; others again have their skulls narrow & small at the base and wide at the top, not unlike the heads of hydrocephalic children. Excessive use of Mercury, inordinate gratification of the venereal appitite, the sudden repulsion of

tanious eruptions, violent arterial action, such
is found in puerperal convulsions, & in fevers
hich attack grofs & plethoric habits, insolation, worms in
alimentary canal, continued attention to any one subject,
e excefsive use of narcotic medicines— Many more might
e enumerated, but these are sufficient to show the
ariety of causes, which produce mental derangement,
some one of its forms.

Madnefs may be divided into three forms, viz;
Furious madnefs, melancholly, ~~in~~ which is considered
hypochondriasis; & madnefs from an imbecility of
ntellect, each of which will require a seperate consideration
this paper. In this, however we shall not attempt those
elicate distinctions which mark the bounderies between
xcentricity of character, & actual madnefs; but con-
sider certain deviations from propriety, as another
egree of the same disease.

Symtoms.

Furious madnefs makes its approach by some peculi-
r symtoms; such as restlefsnefs & watchfulnefs, anxiety
bout imaginary things, incoherent expreffions & ex-
entricity of conversation & conduct, considerable iri-
ability of temper, hostility to friends & relatives, sad-
nefs &, languor. The skin is pale & cool, sometimes the face
s flushed, a singularly wild & vacant stare is visa-
le on the countenence; there is either a total want
f appitite, or there exists a morbidly increased one,
the bowels are obstinately costive, a considerable de-
gree of headach is frequently present. When nearly the
whole of these symtoms are present, we may form a tol-

erable correct prognosis of the character of the ap-
roaching disease. When these symptoms are complete-
y formed, we have a wild, ferocious expression of the
untenance, an appearance of fulness in the ~~[illegible]~~
ace, with red eyes & flushed cheeks, the patient be-
omes very boisterous & clamourous — he whistles
r sings incoherently, makes long & ridiculous speeches,
ses very obscene language, curses & swears most
onably, looses all sense of shame & delicacy, rends
is cloaths, stalks about in a complete state of nu-
ity; and, finally, to finish the mellancholy pic-
ure, the patient degrades himself beneath the
ebute creation, by acting the most disgusting
cenes imaginable.

In this stage of the disease,
he skin is cool, & not the least moisture can be
iscovered on it; and, what is very remarkable,
here is no secretion from the mucus membrane
f the ~~[illegible]~~ nose. It is mentioned by Dr. Rush, &
o satisfy myself of the correctness of the observation,
made an examination of all the patients who were
eranged in the Maryland Hospital, & found it to
e the case in every one, with one exception. The
ulse is sometimes hastened a little, but it is
none frequently natural — The frequency of the
ulse just alluded to, may be referred to nervous
ritation. In this state of madness, one or more
f the nerves of sensation are distroyed, and the oth-
rs become very accute; just what occurs, when a
man looses his sight, the sensation of feeling be-
comes very accute; hence, maniacs hear very accutely

though they taste nothing, and smell nothing.
Different causes, however, produce different symptoms—
Thus the Sarecens, a poor, destitute, & naked horde of
ings; unskilled in the use of arms, or the art of war,
and their imaginations inflamed with a new doctrine,
became partially deranged; & thus fearlessly attacked fortified
wns & citadels; nor did they raise the seige until the
lace capitulated. Here the continued idea, of beau-
ful virgins & plenty of wine, by long action on the
rves of sensation, brought on a disease of the mind,
hich caused them to devastate the most beautiful,
fertile countries of Asia, Europe, & Africa. The same
auses produced the same effect in the wild and
isionary Crusaders, when they marched to the
oly land. The remedies to be used in this disease
epend upon the cause that produced it. I shall,
therefore, say nothing of the treatment until we shall
ave described the different forms of disease.

The next form of mania is melancholly, or as some
uthors call it Hypochondriasis.

The elder physicians were often very unhappy
n their use of terms in medicine. Of this we
have sufficient proof, if we but open their noso-
gical books. Technical words meet our eyes in
very direction, and if they do not confound us,
t least puzzle our wits, in ascertaining their
precise meaning, and original application.

Hypochondriasis is that term which was em-
loyed by the ancient prosologists, to denote an
oneous state of mind, produced, as they conceived

a derangement in the functions of the liver, or some
ther viscus of the hypochondriac region, but, it is, however,
bundantly proven, that the disease is not derived from
diseased liver, or any other of the viscera:— The term is
therefore incorrect, and not applyed with strict propri-
ty.

As this form of mental derangement appears
ost evident in an erronious perception and judge-
ment of things relating to the person & circumstances
f the patient, Dr. Rush proposes to call it Tristamen-
a

Various epithets are applied to this disease in com-
mon conversation, the hypps, spleen, vapours, low-
spirits &c are all synonimous with hypochondria-
is in it various grades. As the author of this essay has
been more in the habit of turning his attention to
the phenomena of diseases than their characteristic
istinction, it will be sufficient to observe that
Dr. Cullen has given the following character of it,
dyspepsia, languor, want of energy, sadness, & fear
rom uncertain causes; with a melancholic temper-
ment."

Persons of nervous & hepatic predispositions
are most subject to melancholly.

Hysteria some-
times resembles melancholly, but there are certain
prominent features, by which they may be distin-
guished from each other. The bloodvessels are
more affected in hysteria than in melancholly,
the nervous system in the latter is always in a state of
torpor, in the former great mobility prevails, hysteria ap-

ars in paroxisms attended with occasional delirium, t is not so with the other — hysteria is alleviated by old, melancholly by warmth.

That it combines with hysteria is proven by many facts. Some melancholic atients are dull & sad at the same moment, at another, they are gay & vivacious to an extreem. They are a-mong the most boisterous laughers, & are addicted to ecat weriment & mischief. These are convulsive effects, nd can only be explained on the supposition that ey are borrowed from hysteria.

Persons advanced in life are more subject to the dis-ase than the young.

Causes

The remote causes operate, first on the body, se-cond on the body through the medium of the mind, & third on both at the same time.

Among the first, are all debilitating occupations of a sendentary kind, fatigue, irregular diet, hunger, watchings, un-wholesome food, chronic fevers &c.

The second variety comprehends all the excesses of the passions & emotions, joy, greif, anger shame, disappointment, vexation, reli-gious destress, intense study &c.

The third kind, or those which operate on the mind and body at the same time, are all perpleaxing employments, where the mind un-usually hurried, & the body fatigued, as professional practice, teaching school &c. the various forms of the venerial disease, by inducing debility, & at the same

ne exciting emotions of shame and sorrow in the mind, ve made many a melancholly patient.

Symtoms.

These are very various. Indigestion, flatulency, costiveness of the most obstinate kind, which requires the most powerful incitants to move the bowels; diarrhea, an excessive secretion of bile, dry skin; & an increased flow of urine, slimy stools; deficiency of appitate; preternatural appitite, cholic, tumed abdomin, tinnitus aurium, throbings of the temples, indistinct vision, palpitations of the heart, burning of the hands & feet &c.

The mind is as variously effected as the body. It is languid, & inactive, and can seldom be employed in thinking on any subject, but that of personal distress. The imagination is in search of doleful images "with a wing that never tires." A thousand fancies are continually excited in the brain; almost every disease lends a share in forming the symtoms of this.

As consumption is the most frequent & fatal disease, it is not to be wondered that it should stand foremost among the chimeras of the patient' brain. If a cough happens to trouble him, he has an abscess in his lungs without doubt; his fears immediately create an hectic fever, should his hands & feet feel a little warmer than usual, it is a certain indecation in his mind, that the ravages of this direful malady are commencing.

Should the unfortunate patient have been lured by a strum-
t, & attacked by the venereal: notwithstanding the poison
been completely eradicated from his system; he is
constant fear of having his palate, or the inside of his
se destroyed, by the inroads of the resuscitated distemper.
his imagination conjures up to his view the loss of repu-
tion & of friends; & predicts ruin & misery through
e remaining part of his life:- Again, he believes
is heart is dilated into an aneurism, or has concre-
ed a polypus, if peradventure through there debility
it palpitates or beats with a little irregularity.
Thus in turns he has a stone in his bladder; an
bscess in his liver; and a tapeworm in his belly.

His mind is effected by certain extraordin-
ry fancies. He is firmly convinced at one crisis
at his body is transformed into a plant & needs the
holesome showers of heaven to make him grow; or,
rhaps, he may imagin himself a teapot: under
is beleif, he is in an awful situation if he be
o roughly handled; and he shivers at the idiea of
eing broken to peices. Probably he has conceived
himself with child; or to sum up the conclu-
ions of his fancies, he persuades himself that he is
expiring, & with no more ado he suffers the pangs of
isolution & lies a lifeless corpse. Nothing, under
these circumstances will quicken him into life,
but some application like the actual cautery—
This powerful stimulant cannot fail to in-
flame his feelings to the most sensible beleif
that he has yet to die.

se are not the worst of the patients mental sufferings.
He is sometimes in despair, here we irresistably exclaim
las poor Yorick!" This state of his mind is induced by a
ll persuasion that he cannot be saved in another world,
d that eternal misery is to be his fate hereafter.
This spiritual distress of the patient, is produced by
wrong views of the character of the Deity, & of his own
ins. Every little neglect in his religeous exercises
- former life, recoils upon his mind, with apprehen-
ions that they were the most fatal remissions of duty, &
he looks forward with all the horror imaginable, to
he time when he will be cast into endless torment-
But it were an endless task to rehearse the almost
numerable symptoms attending this malady. Believing
hat enough has been given to enable the descerning to
form a correct prognosis, we shall now proceed to
he consideration of the last form of mania-

Imbecility of Intellect.

This is either congenital
r adventitious- The first is the effect of some pre-
disposition inherited from the parents; or some mal-
formation of the brain- The second arises from
long continued disease in the system; from sudden
impressions made upon the mind, from grief, & joy,
& several other causes, which it is unnecessary to men-
tion-

Symptoms.

These are generally known, as there
is scarcely a town that has not one or two in it-
They are said to be "foolish;" that they are "cracked"

that they are "hair brained" &c. The patient has an un
meaning countenance; he laughs at every thing & continu-
ally; he makes use of triffling expressions, & is fond of the
society of children, he is irritable in his disposition, care-
less about his dress, or extreemly neat. patients with this
form of madness, are remarkable for their loquacity, they
generally talk when in company, & always mutter when
alone; they are amused with trifles, excessively fond of
sweetmeats &c. hence old people are so fond of their grand
children, that they frequently spend all their time with them.

These symptoms are sufficient to enable us to form a
diagnosis of the nature of the disease, as it is by no means
a difficult one to ascertain.

## Treatment.

We shall hasten on to con-
sider the remedies which are necessary in each form of
the disease— These are "moral & Medical."

When Mania is the consequence of hereditary predisposi-
tion, medicines are seldom if ever of use:— Our treatment
must consist in rendering the situation of the unfortun-
ate patient as comfortable as possible— This may be done in
a variety of ways; by kind & gentle treatment from the
attendants:— no excuse can be received in palliation of an
offence committed against one of those miserable beings—
by keeping the apartments in which they are confined,
well cleansed & ventilated, by using Dr. Rush's tran-
quilizer; thus preventing the patient from rending
his cloaths, & injuring himself— in short, to treat
them just as we would wish ourselves to be treated
if we were in the same melancholly situation.

When it is produced by any other cause, such as in-
uries of the head; here the patient comes under the cog-
izance of surgery & may be relieved by an operation—
hen from intoxication, we must abstract the stimulous
lowly & administer emetics & cathartics, anodynes of
ome narcotic, opium, perhaps is the best; from in-
lation, the antiphlogistic treatment must be resort-
d to, cold applications to the head; the strictest regard
must be had to his diet, he should be kept in a room
here he would not be exposed to two much light; his
owels should be kept solu~~ble~~, with neutral salts &c.
When from onanism the patient must be confined in
uch a manner, as will prevent his continuing so shame-
ul a practice, when from excessive venery, he must,
n the same manner be prevented from indulging
is sensual appitite— The man of feeling, may, thus
ery frequently alleviate the symptoms if he can not
emove them. Labour has been recommended by some
riters; but in that form of mania, to which the above
treatment is applicable, it can seldom be resorted to, as
he patient is in such a state, that he cannot be managed
when at liberty.

The above treatment is applicable
in that form of the disease, which has been called fu-
rious madness.

We shall now go on to consider the treat-
ent in the second form of the disease, or that which we
ave called melancholly.

The remedies for this, form them-
elves into two kinds: viz. those that act through the me-
ium of the alimentary canal, & those that produce

eir peculiar effects on the organs of sensation —

When the disease is attended with symptoms of in-
gestion, costiveness, flatulency, diarrhea &c. as al-
ready enumerated, medicines which relieve those sym-
ms must be given — Indigestion must be obviated
a strict attention to diet, costiveness must be pre-
ented by laxative medicines, & diarrhea by correcting
e morbid secretions of the intestines. An excessive
cretion of bile should be prevented by alterative med-
ines, such, as small doses of calomel, the blue pill, or
e nitro-muriatic bath, exercise of various kinds may
resorted to, such as riding on horse back, or in a
riage, swinging & sailing; moderate exercise on foot
as been recommended. Thus, the judicious practi-
oner, by a proper regard to symptoms, will generally
sure success.

With respect to the remedies which act on
e organs of sensation; these are such as may occur to the
actitioner at the time. The disease occurs, as has already
een stated under a variety of forms, & our treatment must
e regulated accordingly: - Thus, if the patient imagin him-
elf some great man, we must favour the belief, & treat
im, with all the care & attention, as if his disease were a real
e; & observe all the respect, in his presence, which such charac-
rs are in the habit of receiving, until an opportunity may
resent itself to dispell his visionary greatness. If he im-
gin himself transformed into an animal, as is very fre-
uently the case; such as a dog or a cat, or even a hog, he
ay have his food thrown to him in the same manner
s we would feed those animals; ~~we may~~ by this ce-

131

An
Inaugural Dissertation
on the
Yellow Fever.
Submitted to the Examination
of the
Provost, Medical Professors, and Trustees,
of the
University of Maryland,
for the
Degree of Doctor of Medicine,
on the seventh day of April A.D. 1838.
by
James R. Ward,
of Maryland.

To

Nathaniel Potter. M D.

Professor of the Theory & Practice of Medicine.

in the

University of Maryland.

This Dissertation is humbly inscribed, as a tribute of respect, for the great kindness and attention received from him; and which, will always be remembered, with gratitude by

his much indebted,

and affectionate Pupil.

The Author.

# A Dissertation &c



From the prevalence of this disease in the United States, & from the mortality which has attended its appearance on our sea port towns, the attention of Medical men was generally attracted. The most distinguished of the age, in the profession of medicine made every exertion to discover the cause, the seat, & the best plan of treatment.

Great credit is due to professor Rush, for the first correct theory of the Yellow fever. Its pathology was soon determined from Post mortem examination, accurate observation, & experience. No disease perhaps has had more talent engaged in its examination than the present one; & none perhaps has caused so much litigation; and still one or two points are sub judice. I refer particularly to the doctrine of contagion. The physicians of the North, firmly believe it to be contagious; & some deny that it ever originated in this country, but that it is always imported.

In writing an inaugural dissertation on this subject I am fully aware of the ability, with which it has already been handled. by men who will be a lasting honor to the profession, & whose usefulness will not cease with their existence. This rendes me more diffident in writing on this subject, however, having seen the disease in in a climate that is not generally known & whose diseases have not been discribed. I thought perhaps that some remarks might be made, which would throw some light on this important & very interesting subject. If this feeble attempt should be useful in any way to the science of Medicine, my exertions will be amply rewarded

Authors in describing this disease have given it various names. The Yellow fever of the West Indies is the most common for this disease, but as yellowness is one of its symtoms & which can generally be prevented by proper treatment, it certainly cannot be properly used. Dr Rush calls it the Bilious remitting Yellow fever, which, is certainly a more appropriate name, & one expressive in some measure of the nature of the disease. It has also been called the plague, however I have adopted the common term —

We have says professor Rush, this fever, like the steady blowing of the equinoctial gales, we have it again assuming a milder remittent type, sometimes imitating the character of common intermittent fever, but rapidly paving its insidious way to the distruction of the patient. From its ever varying type, for says Professor Potter, we find no two epidemics requiring the same treatment, & wearing the same livery. It will be admitted that the disease as called by Dr Rush is the most appropriate it has yet received.. If another should be given, any word that would define accurately the appearance & expression of the eye, would be the most appropriate. Language however it will be found inadequate to describe it, yet once seen, it will never be forgotten.

## Causes.

This is another point which is not settled, many Physicians of the present believe the disease is always imported, & hence the strictness observed, in some of our sea ports as, regards the Quarentine Laws, which are a disgrace to all concerned in their enactment, for they carry on their very face, the strongest marks of folly & ignorance. There are so many facts, that prove beyond controversy, that the disease

has originated in places that have had no connection with foreigners, or vessels from places where the disease prevailed. Marsh Effluvia is generally admitted, among those, who have seen the disease frequently, to be the cause, not only of the Yellow fever but also of the common bilious & intermitting fevers of our country; as a necessary consequence they would be deemed by those who believe it is a foreign disease. Let us examine into the facts which present themselves, to those who reside in that part of the country where these fevers are prevalent. On the Eastern Shore of Maryland, where the bilious and intermittent fevers are common every year — Foreigners who visit these countries, during the summer & early in the fall are seized with the bilious & yellow fever, & die with the black vomit, which those who are opposed to this doctrine consider as placing the character of the disease beyond doubt, whilst those, who have been raised & continue to reside there, will enjoy good health, or have the ague & fever, which they do not regard. Now the same cause is acting on both at the same time, & producing diseases, of the same kind only more violent in degree. Stronger evidence could not be asked, than the diseases of the American Colony of blacks at Cape Mesurade on the coast of Africa. These people when they first settled on that coast, died in great numbers with the Yellow fever, so great was the mortality, that it was thought it would be necessary to abandon the settlement, they have however become acclimated, & now enjoy good health & what are the diseases to which they are now most subject? Common, mild intermittents; when this fever is prevalent, what do we find among foreigners, who visit that place? a violent & fatal fever, which destroys them in two or three days. With the black vomit the last struggle comes on, & death soon closes the scene. Let one of these blacks return to the United States for a winter

and then return again to Africa & he will have the Yellow fever. This has been repeatedly observed in the inhabitants of the West Indies. for, if after residing immediately on the shore, or in the large towns, they go into the interior of the islands and remain for some time, upon coming down on the coast, they will be seized with the Yellow fever.

This is proof sufficient to convince the unprejudiced mind, that the Yellow fever & Bilious fever have a common cause & that they differ only in degree. To account for this difference certainly cannot be difficult. The habitual Laudanum taker will, with impunity receive at one draught, a dose sufficient to destroy two persons who are unaccustomed to its use, without any inconvenience to himself; in the same way the Constitution becomes accustomed to the action of this poison, so much so, that it may not produce any disease, but the atmosphere that one man breathes with impunity, who has been habituated to it, will, soon produce violent disease & death in him who had never been accustomed to its action.

The Yellow fever as described by Professors Rush & Potter, as it occured in the United States, at or soon after they wrote, bore the same general character as that I was witness to, in the West Indies & required the same active treatment. I had an opportunity of feeling that state of the skin, which is produced by Mercury & which Dr Good calls a "diaphnoe", this was the first evidence I had of the Mercury affecting the system. From Dr Goods accurate description of this state of the skin, I remarked as soon as I felt the patient that the mercury had had the desired effect, which was manifested in a few hours from the patients complaining of his mouth. It is not requisite that I should describe the Yellow fever of this country which has been so ably done, by men whose celebrity will be the proud incentives to acc-

tion in the youthful candidate for Medical distinction, these remarks I thought necessary to make before attempting to describe the disease as it occured on board the United States Schooner Shark during a cruise of six weeks on the coast of Africa. The Shark sailed from the U. S. in November & arrived at the American Colony at Cape Mesurado in February. some few of the Colonists were labouring under chills & fever, generally they were healthy, our crew was remarkably so, when we arrived & continued so for some times, as we generally were at sea every day or two & the men were not much exposed. Before describing this disease, as it occured on board the Shark. I think it necessary, in order to elucidate the subject as much as possible, to make a few remarks on the country & climate. these cannot be as accurate as I could wish, but shall be as perfect as my limited observation atended. The Country is thickly covered with low wood, and large timber. The soil is very rich, none perhaps more so: none, where vegetation is more rapid, & luxuriant. The seasons are divided into wet & dry. During the wet season, the rain falls in torrents & almost deluges the face of the Country after a heavy shower the air is close, & together with the perpendicular rays of the sun renders it almost insupportable. When the dry season comes on vegetation thrives astonishingly it soon grows rank, fall, & putrefaction takes place; Large collections of stangnant water filled with vegetable matter in a state of putrefaction, are to be found along the banks of the river. We have the diseases arising from Marsh Effluvia prevailing during the year as putrefaction is continually going on. Yet they are more violent at one time of the year than at another Our stay was during the dry season. In the morning until the sea

breeze sets in the air is warm & oppressive; about 9 O clock we have the sea breeze which is refreshing & invigorating. In the afternoon we have the land breeze which produces a change in the temperature of the air, that is truly surprising. It becomes cold, chilly, & damp, & warm clothing is necessary. Many degrees difference will be found in the Thermometer during the twenty four hours. In the afternoon the mist is always so great that the sun becomes obscured about 4 O clock.

It may easily be conceived that the air coming over such a country, would be loaded with moisture & the poisonous effluvia which is continually evolved from these wide extended marshes.

What Constitution could stand with these changes occurring every day. Cold, the great enemy of our race, here acts with certainty; predisposing the body to receive the disease, the cause of which disease is continually acting & surrounding him. his anxiety & dread which a knowledge of the unhealthiness of the climate increases, also renders him, a more easy & certain victim to his insidious foe.

All those who were taken with this fever complained of violent pain in the head & back, great uneasiness in the Epigastric region, & more or less irritability of the Stomach. all had the red, muddy & glassy eye. From the want of the written account of the most violent case that occured; I am unable to be as accurate as I could wish. The patient was a stout man & of good constitution, he was seized with a violent chill, soon after rising in the morning, succeeded by a high fever, pain in the head and back great irritability of the Stomach & great uneasiness at the epigastric region —

The state of the pulse, I do not, correctly remember, but to the best of my recollection, it was depressed. he was bled freely that night & took Calomel during the day, no evacuation from the bowels took place, & an injection was administered. the next day the patient was able to walk on the deck, & complained but little, every thing looked favourable, except the eye which still continued to have the same unnatural appearance as at first. On the third day in the morning, he was again seized with a chill, followed by fever, in the afternoon he had another chill. On every other day he was better, the patient not appearing to be much debilitated for the first 2 or 3 days. the active treatment was continued, as the evacuations shewed that the liver was deeply involved; for notwithstanding the continual vomiting no bile was thrown up from the Stomach. On the fifth day the debility became very apparent & his countenance indicated much distress. it was now found necessary to change the treatment. Tonics & stimulants were ordered. On the morning of the sixth he was evidently sinking, about 4 in the afternoon he was seized with convulsions, followed by the black vomit & delirium. the patient on being asked how he felt answered, horribly! at 8 he was again seized with convulsions & expired with the black vomit running out of his mouth. Yellowness here did not take place to any degree until after death, when the whole surface became yellow. this was the first case that occured & the only one that terminated fatally. To another who was seized with same symtoms a large dose of Calomel was administered, & the patient was sponged

with vinegar and water, during the continuance of the fever to great advantage. Tonics were given as soon as possible & the patient soon convalesced.

The sponging has been much recommended, & it certainly has been found highly useful. It is hazardous, in fact impossible to pursue the same active treatment as in the diseases of this Country & the West Indies, as experience warrants me in asserting. On our passage to the United States from Havana several of our men were seized with the Yellow fever, the symptoms were not so strongly characterstic of yellow fever, as those on the coast of Africa. but yet required several bleedings & active cathartics, & the fever was not removed until some mercurial impression had been made on the system. This fact shows the difference between the diseases of the two countries. One of the Colonists who has paid attention to the disease, after considerable observation remarked, that he found that foreigners would not bear an active treatment more than a day or two & then required to be supported. The most successful treatment of the disease was, to use strong Mercurial cathartics in the first instance venesection being cautiously used, together with sponging & diaphoretics. They have an article indigenous to the country which is found to be highly useful, as a diaphoretic & also a decoction of the leaves of the orange & lime tree is also useful.

The early use of tonics is absolutely necessary as they otherwise soon sink. Under this plan of treatment the disease is found manageable and a voyage to sea soon restores your Patient.

Connected with this subject, there are two circumstances of some importance, the one is that remaining on shore all night is almost always attended with sickness. Out of 4 who remained on shore all night, 3 were sick and one died.

A physician who resided there for some time informed me, that after he got well enough to go on board ship to sleep if he left the shore before sun set, he escaped his chill but he invariably had it if he remained for any time after that period. Johnson on the diseases of tropical climates has also mentioned a fact of this kind. The other is, that constipation for two or three days is followed by high fever & is often fatal to the foreigner, whereas those who reside there feel no inconvenience whatever, this say some is easily explained, the liver is in a state of congestion & the intestines have not their accustomed stimulants from there being no bile secreted. If this be the cause, ought a cathartic, which only evacuates the contents of the alimentary canal relieve the liver and restore the secretion. I am of opinion that it may as a congestion might be produced in the liver which has already been debilitated by the action of heat & consequently and ~~consequently~~ more sensible to the effects of Marsh Effluvia, by the operation of an active cathartic. What is the effect of a warm climate on this viscus? it stimulates the organ to action & we find the quantity of bile increased: the long continued application of stimulants to any organ will debilitate it, heat as the writers on warm climates prove has a stimulant effect upon the liver. It consequently produces debility of the organ, & it becomes more liable to disease, Marsh Miasma acts on it with power & the result is disease. The faeculent contents of the alimentary

I believe him and if there wanted other thing to shew that the Origine of Bilious fever was the same ~~as~~ that the ~~same~~ materials were the same (only varying in Concentration)

Canal, I should suppose must have some effect on fevers of this kind. The Physicians of the West Indies endeavor to restore the secretion of the liver by the use of Calomel in large doses, but they also pay great attention to the removal of the faeces, & keep the bowels soluble. My attention was called to this subject from a case of typhus fever, under the care of Professor Rush & mentioned by Professor Potter in his lectures. This patient had been stimulated for some time, Dr Rush raised him to a certain degree by tonics &c, but he became stationary, notwithstanding the most powerful tonics were used, a diarrhoea seized him & his convalescence was rapid & the use of tonics was no longer necessary; after having heard of this case. I saw a patient who had a low muttering delirium, stupor & difficult to be roused & scarcely answering the questions that were put to him.
Yet all these symtoms were removed by the free action of a cathartic, which had been given whilst the patient was labouring under these symtoms and the pulse which was before weak & small became fuller & softer & the patient fell tranquilly to sleep.
I think if more attention were paid to the bowels in these fevers, our attention would be rewarded by the speedier convalescence of our patients. Let it not be understood, that I approve of purgings in all stages of fever. If I did, patients in certain stages of fever, would soon be hurled into Eternity. From the sympathy that is observed to exist, between the bowels & various other organs in different diseases, I am induced to believe, that a soluble state of the bowels would be more advantageous than is generally supposed, & that we are oftener deterred from this practice,

time commencial, holdeng it to be a fact Mathemat-
icolly true where ever the Cause grows, there also
may be seen the effects — and the Apetites of Dr.
Warren that the disease was imported into Charlestown in
one of his Majestys ships from Palestine ~~is all~~ in 1721/2
is all a farce, we say that in those islands were found
all the materials requisite to the formation of this disease
that there the Thermometer ranged above ° there
also could be seen Animal & Vegetable putrefaction
with the requisite proportion of moisture — there
was an excessive race of Men and so when we turned
our eyes from the Cause we beheld the effect, the account
given by writers of the day and the account recieved from the
the words of the Aborigines Them selves, who the having no
Idea of this disease, and Nothing to guide them but the
Wild and uncultivated eye declare that when one
would die of these great rechuffs he would exhibit
that colour so peculiar as to the yellow fever above —
Dr. Mosely says the yellow fever is the disease noticed by on a specimen of the
Inspirators and the head Kausos — Also noticed by Aretus
and Galen by the term Febris Ardens or Causus. the In-
fluence of Climate & season on diseases are too obvious on
the Character of diseases to be denied. and a learned
author tells us this disease is widely different as it appears
with West India Islands from that which appears in
the mild and temperate climates of Europe. and tells
us in the mild & pleasant Countries of Europe they have
the Bilious Remittent fever in a very gentle form — I believe

from our too great fear of debility. The prophylaxis is the same on the coast of Africa, as advised by Dr Potter & others as regards the disease in the United States. The bowels to be kept soluble, diet moderate, abstaining from ardent spirits unless accustomed to their use, carefully guarding against exposure in the sun, & to the night air; keeping the feet & body dry, never remaining on shore after sunset, & by no means all night, a careful attention to these directions, will prevent the disease or render it lights. As to the question of its contagious character I am decidedly an ante contagionist. Were these diseases contagious, we should have had none left to tell the dismal story of our miserable end, as ten persons in one case slept in one apartment with a man who died of the disease, & the marine guard who were the only sufferers in the West Indies, from being exposed at night on post. were confined during the whole of their sickness, on the same small deck where 20 or 30 sailors slept every night, & they slept almost in contact with each other, & yet not one had the disease, for my own part I visited these men frequently, & never had the least fear on account of its being contagious.

These remarks have been made with diffidence. To be an author so young, was far from my wish. Yet necessity hath compelled me to the task. The inaccuracies of youth must be looked over, when improvement may be expected from the stimulus to exertion that is given to the youthful enquiring mind, from reading the writings of a Rush & many others, who have done honor to their age, & country, & last, though not the

to Marseilles, and thence to Martinique about the year 1721 or 2 — Mr Hughes tells us that Dr Gamble remembers the year of 1691 that a fever prevailed in Barbados called pestilential fever new distemper and Bilious fever and it was most fatal in May June July & August — Martinico appears to have been the place which this disease first made its appearance It would be impossible for us to say when or where this disease first made its appearance, we find from accounts of very early dates both from Europe and amongst the Aborigines of this Country that prevailed at times fatal epidemics in whose Countenance we be held ~~the~~ some of the features of this disease sweeping away those unfortunate souls exposed to its power, but though we have not had sufficient documents from the dark and Obscure ages of Antiquity to say positively that it was this disease which destroyed the inhabitants of the west India islands in 1721 & 22 and in many other places yet it is possible ~~where ever the requisite was~~ ~~is what was~~ it might have been for aught we can say knowing all the materials requisite for ~~this disease~~ formation of this disease who present in those Islands and many other places and from our present knowledge the Etiology or the Causes of the disease, we believe it has escaped

least the deserved celebrity of the professors of the University of Maryland which has been & will continue to be the proud Alma Mater of many worthies of our country & profession —

Finis.

When this disease first sprung up in the western Ho
it was considered of the same species as the Asiatic
plague, and only one disease varying in degree — it
called by them the plague — some time after
this it was noticed under the head of Yellow fever
or Black Vomit derived its origine from two of the
most prominent symptoms in the disease
The Ancient Authors speak of a black and
Bilious Vomiting but not as a symptom of
importance but rather as we have yet
received notice of the great diagnostic the yel
low Colour Dr. Hodges mentions One case
in the plague where a man turned green — and
from numerous circumstances we are compell
Comparatively speaking to call yellow fever
disease of Modern Origine — Dr. Hodges says Dr. S
Hellan as we have no account of this disease
from the Ancients who lived and practiced
hot climates we must give it some name
and he gave it a name a much more ap
able one than any Author in his time the
Bilious pectic fever — and says from the
best from the best of his knowledge it is a
disease indigenous to hot Countries — particu
by the W Ind islands & the Continent of America
— Dr. Warren thought that this
disease was originally brought from Africa

157

# On Phthisis Pulmonalis

An

Inaugural Dissertation

submitted to an Examination

of the Faculty

of the University of Maryland.

for the degree of Doctor

of Medicine

by

Horatio G. Grieves.

—1828—

To

Doctor Frederick Dorsey

of Hagers-town,

those sheets

are most respectfully inscribed

by his friend, and pupil,

H. G. Grieves.

To

John M. Lawrence M.D.
of Cumberland

Knowing of no individual who unites in his ~~talent~~ character more talent and benevolence than yourself, permit me to inscribe to you, this "stumbling block" in Medicine, as a token of the sincere respect and estum entertained for you by the

Author

To

John M. Howland M.D.

of Baltimore

There is no individual, within the circle of my acquaintance, to whom I could with so much pleasure, or so much propriety, dedicate the contents of the following sheets, as to yourself. For that disinterested solicitude, manifested toward me whilst pursuing my Medical studies under your able instruction; accept the gratitude you so justly deserve. That the propitious smiles of heaven, to which you so yealously aspire, may ever be yours is the sincere desire of

Your friend, and pupil,

H. G. Fievs.

# Dissertation upon Phthisis Pulmonalis.

——————— Study well the clime?
Mould to its manners our congenial frames,
And mitigate those ills, we cannot shun.

The disease upon which it becomes us to treat has been emphatically as well as justly ranked among the "aspprobria medicorum"; and as an incontrovertable evidence of this fact our bills of mortality are a lamentable testamony: to use the words of an immortal sage whose time and talent were studeously, zealously, and indefatigably directed to the advancement of his profession in a degree almost bordering upon enthusiasm — "Phthisis Pulmonalis as it is justly denominated is of all diseases one of the most fatal." When we take a circumspective view over the extended field of nature and as idle spectators animadvert upon our insignificance and imbecility; what description of

subjects do we find fall victims to its deadly influence? Alas alas! but amongst them stand conspicuous those who have arrived at the most engaging period of life and those perhaps the fairest of our race. Must talent then be spent in vain? Must researches then be spent in vain? That same enterprize which snatched lightning from the clouds. That same philosophy and erudition, which has thus high elevated a scientific profession; it yet remains for to discover a specific for what we conceive to be the greatest enemy to the human race.

Phthisis Pulmonalis is understood to be that affection in which a general wasting or consumption of the body arises from a diseased of the lungs.

Notwithstanding the want of Information upon this disease, which has been manifested by some of the most enlightened of the Profession much

diversity of opinion as regards its Pathology has been advanced. Whilst by some it has been doubted whether the disease strictly speaking can be considered Idiopathic. By others it is considered to be invariably symptomatic of other diseases. Pneumonia - Catarrh - Scrofula - Haemptesis - and several other genera have been mentioned as being the exciting causes of Phthisis; but this with the rest of the Pathology must yet remain a desideratum. The theory however of the disease being one "sui generis" appears in itself reasonable and we find it substantiated from the circumstance of the disease making its appearance and approaches in such an insidious and formidable manner as frequently to be beyond the reach of human skill before its deadly effects shall have been promulgated.

It has been remarked that a peculiar description of subjects are more susceptible

of this disease than others and those have been described as being of a scrofulous diathesis. Whether any real advantage is to be derived from this kind of distinction or not it yet remains to be discovered.

Phthisis Pulmonalis has as an attendant in every instance that modification of fever termed "Hectic" this great and prominent characteristic has been described as making its appearance sometime before a fatal termination takes place - much controversy of opinion has been entertained as regards this of all most distressing symptoms it therefore becomes us to remain silent untill further and more satisfactory investigation be reported.

Writers when describing disease have thought proper to adopt by way of perspicuity certain classifications, and in order that our present object may be facilitated we for similar causes will adopt a plan approved; although disapproved

of by some upon whom the meed of praise has not been lavished undeservedly. The causes and symptoms of Phthisis have been modified and magnified to so great an extent by nearly every one who has attempted the subject; that it is less a source of astonishment than of mortified feeling, why so much yet remains undiscovered. We will now proceed to a description by observing an arrangement which to us appears plausable as well systematic.

Those species have been denominated the Catarrhal, the Apostematous, and the Tuberculous, In every modification of Phthisis, the symptoms have been referred to three heads — the Pneumonic or Pulmonary symptoms the symptoms of Hectic Fever & the supervening or what has been termed consequent symptoms. We will briefly commence by describing the Pneumonic symptoms as occurring Catarrhal Phthisis

Cold is looked upto as being the great fountain and throne of all Catarrhal diseases; this modification of Phthisis makes its approach with that same undeviating train of symptoms which characterize this class of disease. After the more violent Pneumonic symptoms shall have subsided, others of a more distressing character are said to develope themselves, and first "Tussis" which may be described as being one of the most prominent features of incipient Phthisis this peculiar disagreeable and troublesome cough we find much aggravated when in the recumbent posture, at the same time we do not find it wanting for disagreeable sensation while in any other position. After this the chieftan of the van shall have proceeded with an almost irreparable devastation, others, of an equally alarming character follow up the afflicting scene.

Dyspnoea - sense of soreness in the Thoracic region -

and with these attend an expectoration which by this time assumes a yellow aspect, disagreeable both in taste, and smell: occasionally instances we find mingled with the matter expectorated streaks of blood said by discerning men to be an indication of a scrofulous diathesis. Upon this latter clause it remains not for the inexperienced to animadvert, suffice it to say that the medical opinions of medical men should always be open to impartial investigation as well as rigid scrutiny and if erroneous or inconsistant with Medical Philosophy talent, and experience, and aspiring ambition, it becomes to controvert. We regret that it is not in our power to dwell more extensively upon this as we conceive to be important point; the confined limits of an Inaugural Dissertation, together with the reflection, of its perchance prove uninteresting to the gentleman into whose hands those sheets are consideration abundantly obvious

to justify us in abandoning our contemplated arrangement.

Another variety has been described as being the Pneumonic Symptoms occurring in Apostematous Phthisis. This variety has been described also as being much less frequent than former, its symptoms more prominently characterized and its result a more fruitful source of lamentable contemplation. Apostematous Phthisis is a consequence upon a large abscess formed within the cavity of the Thorax; the former has been described as being ushered in with cough, and other catarrhal symptoms and as being a consequence upon atmospheric vicissitudes: whilst the Apostematous or the stage we are about to describe would be by those who advocate the theory of the disease being a con-

sequence upon certain other diseases and by those who mingle causes with effects attributed in all probability to an Pneumonia, to an Hemoptises, or to some other Inflamitory thoracic disease; we do not conceive that any important advantage can result from this kind of distinction, but we do conceive it to be of vital importance to know whether certain diseases as they are described to be are exciting causes of Phthesis Pulmonalis or what plausability would make them appear to be merely precursory symptoms. Upon this point much of a speculating nature has been invested in, which has not only had a tendancy to render ~~intricate~~ what might otherwise have been considered simple but to render still more complicated a complicated subject.

Authors accounting for abscess forming in the chest af-

signs as its most frequent cause an haemorrhage taking place from the lungs. Dr Cullen has dwelt upon this subject with so much pertinacity, that he has ventured to rank it under the genus of Haemoptysis, and as is generally presumed erroneously viewed it a sequelae of that affection: that this variety of Phthesis Pulmonalis does often commence with a spitting of blood no one pretends to deny; but it would be equally rational and equally justifiable to place Intermittent under the genus Cholera, because it has as a frequent precursory symptom Nausea and Vomiting. Vomiting may then be considered as a consequence of Haemoptysis or spitting of blood from Pulmonary Predisposition. Blood though emanating from the lungs is not to be considered as the only source

of this variety of disease, Inflamation of the Lungs as well as external thoracic injuries deserve a conspicuous part in this of all most afflicting tragedies. Pain, Dyspnoea, and other Phthisical symptoms attendant upon the incipient stage of Apostema are said to be comparatively speaking inconsiderable, in this state of things severe fits of coughing will frequently supervene attended with but little expectoration; and in this point consists the great diagnosis between catarrhal and apostematous Phthisis. We had intended dwelling more at length upon this variety of the disease, but when we find Philosophical sages here drop their pen, surely it must be an indication of their having reached a "Ne plus ultra" point.

a point not to be transcended by the inexperienced however enthusiastic. We come now to a description of the most frequent as well as most dangerous of all the species of Phthisis.

The Pneumonic symptoms attendant upon the Tuberculous Phthisis approach in many respects the variety last described it would therefore be superfluous to recapitulate what has already been said. Tuberculous Phthisis in consequence of its arising from the greatest variety of exciting causes, is of course of most frequent occurrence. But however numerous and varied those exciting may be all of them operate as giving rise to this variety or modification of Phthisis, on one general principal,

viz. as inducing those peculiar tumours termed tubercles which have been found in the lungs of patients who have died in this stages of Phthisis: the symptoms attendant upon this variety of the disease are comparatively speaking inconsiderable, in the incipient stage cough and other Pneumonic symptoms are by no means urgent, and in most cases no remarkable pain of breast attends, and when it does it is neither fixed or constant; and contrary to what happens in Apostematous Phthisis the patient can lie with equal facility on either side. Most frequently the matter expectorated in this stage of the disease is a thin watery fluid slightly tinged with

blood, it has been described as having very much that appearance of sanies which is often discharged from scrofulous sores. When this state of expectoration takes place "Hectic fever" is seldom wanting to a considerable degree.

Much controversy of opinion as regards the pathology of "Hectic" has been entertained by some of the most enlightened of the profession. After viewing strictly the different theories advanced upon this subject we are inclined to consider that which arises from a certain irritable state of the system induced by an absorption of pus to be the most probable as well as plausable conjecture.

"The Hectic Fever" is very generally not of the continued form - but consisting of remissions and exacerbations; and from this circumstance it has been justly described as being a variety of Intermittent. All that is necessary to mention in this state of things is the description of Pulse attendant upon hectic fever, which in most instances is seldom observed to be under 100 and frequently above 120 strokes in the minute. In some instances of Hectic the pulse is full and hard; but more frequently small and hard during the paroxism as well as intermission. After this distressing symptom shall have been observed for some time; others

of an equally alarming character make their appearance, those have been termed the superveming and then again they have been considered as sequelae of "Hectic." It is doubtful in our minds whether they can be considered strictly speaking either; they may be perhaps like hectic symptomatic of the progress of the disease from an absorption of ill-conditioned matter from the Lungs, the probability is though that if not induced they are at least much aggravated by the hectic fever.

A picture in itself deplorable is now we approach with a feeling no less painfull than a consciousness of our inadequacy to depict.

Human misery is in no instance more strikingly exemplified than in the one which now presents itself to our consideration, and the one which now it is our province to describe. After all the foregoing distressing symptoms shall have made their appearance upon the stage of humanity, characters equally conspicuous terminate the tragical scene. Emaciation or wasting, has been justly numbered amongst the prominent Pathognomonics, of confirmed Phthisis Pulmonalis, and in no part of the system is this symptom more obvious than in the face, here, it has been termed the "Facies Hippocratis" in short the whole system becomes so completely reduced that it has been represented

to be as it were a living skeleton. With emaciation attends aphthous spots both upon the tongue and mouth, which in the end not unfrequently degenerate into ill conditioned ulcers. These aphthae & consequent ulcerations are particularly observed in the apostematous phthisis, when the expectoration of purulent is considerable, and when, in place of the appearance of mild pus, the discharge has degenerated into an ichorous state. This affection affection of the mouth has by some been ascribed, and probably not without reason, to the action of the expectorated on the mouth. To the same cause, acting upon other parts of the alimentary canal, especially

on the intestines, has been ascribed another symptom, very common and very distressing, the colliquative diarrhoea, as well as colliquative diarrhoea colliquative sweats, With these as they are called supervening symptoms, and the continuance of Hectic it is by no means surprising, that the disease after arriving at this state should in almost every case have a fatal termination. Having thus superficially mentioned the causes, and symptoms of Phthisis we will now proceed to a part of our subject, which we promise shall neither be tedious or lengthy

Much has been said about the treatment of this disease, and as in every other disease it is our provence to remove as far as is practi-

cable and as soon as is practicable, all exciting causes; and thus it must be an object with the practitioner, to counteract, as far as is practicable the ill-consequences of purulent absorption. The means of affecting this removal must, in the different species, be accommodated, to the nature of that particular source from whence the purulence is furnished. In the Catarrhal Phthisis the source of that purulent matter which produces the symptoms is a mere inflamed surface; hence, our first indication of cure is to diminish that impetus with which the blood passes through the systemic but more particularly

the pulmonic circulation, a diminution of this impetus may be obtained in different ways. The second indication requiring to be fulfilled in the cure of this variety of the disease, is the restoration of a natural condition to the vessels of the lungs, and with a view of fulfilling this indication, much more is to be derived from regimen than from medicines, and particularly from gentle exercise, pure air, and a mild nutricious diet. In the treatment of Apostematous Phthisis much cannot be said, and we regret much that the same words are applicable, to the third and last or that stage termed Tuberculous Phthisis. Near-

by all of the most eminent writers, have introduced articles in the treatment of Phthisis Pulmonalis described by them as specifics in the disease, we must content ourselves by briefly mentioning those which have mostly excited the attention of the medical world.

We had occasion before to remark that different means had been employed for the purpose of lessening the force of the heart and arteries, and for this purpose the Digitalis of Dr. Beddoes has not held an obscure rank, but to speak in the words of an illustrious Potter like its author has sunk, perhaps to rise no more, this remedy we conceive to be not

only dangerous in its effects, but inefficient in its consequences, it should therefore deservedly be abandoned from the catalogue of medicines employed in the treatment of Phthisis Pulmonalis. Another article the deleterious effects of which upon the animal economy stands unrivalled within the covered of materia Medica, has excited attention in no small degree from the votaries of the healing art; and notwithstanding the train of pernicious symptoms, necessarilly consequent upon its administration, sages worn out in the shafts of professional prosecution, sages whose eminence

and whose immortality has strengthened with their strength, and grown with their growth; have placed it as a curative in a superlative degree.

M. Gay Lussac says he has constantly found that the Hydrocyanic acid administered to the unhappy consumptive, not only appeased pain, but diminished frequency of cough, moderated and rendered more easy expectoration, and lastly procured the patient sleep at night, without any colliquative sweats; he goes on still further to state, that those who are accostomed to follow the march & progress of Phthisis and witness the suffering without number of those individuals attacked

by this terrible malady and by which they are overpowered, will easily appreciate the benefit of this practice. Encomiums equally extravigant have been lavished upon the article, by men equally eminent, and perhaps equally aspiring. With regard to the employment of Emetics in this disease: they have been mentioned as being in all probability servisable, when not given in large or frequent repeated doses, of this class of medicines Tart: Ant: has been selected as being most servisable. All the class of Demulcents have been employed and as palliatives are considered to be very appropriate remedies. Another important remedy yet remains to be

treated of; the administration of Calomel, particularly in the inflamitory form of the disease, which we view as being the only form that can be cured, has acquired celebrity in a degree of no small consideration. Bloodletting in conjunction with the last mentioned remedy appear to be means placed in our hands by which we are to grasp from inevitable destruction myriods of the human family.

With those crude and undigested remarks we close, soliciting from the generous reader a liberal perusal of the same, and reminding him that upon our part it has been a matter of compulsion and not of choice.

217

An

Inaugural Dissertation

on

Cynanche Trachealis

Submitted to an examination of

the faculty of the University of

Maryland

for the degree of Doctor of Medicine

by

John O. Wharton A.M.

of

Tennessee.

1828

To

Nathaniel Potter M. D.

This Juvenile Production is

most respectfully inscribed

By

The Author—

This is a disease apparently of Modern origin. I say apparently because altho it may have existed yet it has not been described by any of the Ancient with even a tolerable degree of accuracy. To Dr Home of Edinburgh is justly due the credit of having first called the attention of the Profession to this interesting subject. Dr Cheyne has given the fullest and best account of this disease. Since the enquiry of Dr Home into the "Croup" it has been noticed by many Authors, most of whom have at least given a different appellation to the disease, if

they have advanced nothing new with regard to its Pathology. Dr Scott of Edinburgh has called it Asthma Acuta, from the difficulty of Breathing which constantly attends the disease. By Dr Russel it has been termed Angina Inflamatoria. A variety of other names have been given to it; By one, it is called Catarrhus Suffocativus by another, Morbus Strangulatorius and by a third, Cynanche Stridula But of all its names that which has been adopted by Profr Potter seems to me most appropriate. Tracheitis

The Croup mostly attacks children between the second and seventh years of their age: a complete immunity from the disease is however enjoyed by no age. Adults are much less liable to attacks of this formidable malady than infants. This seems to be owing "says Dr Gregory" to some alteration which the mucous membrane of the Trachea undergoes about the age of Puberty. Altho the disease generally appears in a sporadic form, yet it sometimes

prevails as an epidemic, as was lamentably case in Virginia in the year 1799, when the Father of our country fell a victim to it.

It is not my intention in this essay to inquire concerning the great diversity of opinion which exists among Medical men, with regard to the Pathology of the Disease. Tracheitis or Croup is an inflammation of the Mucous Membrane lining the Trachea, from which it is often extended to the muscles surrounding the Glottis; the case has over it sometimes reversed, the disease commencing in the Muscles, and travelling downwards.

In the latter instance it must be confessed that some degree of spasmodic contraction of the muscles will occur; such however is the situation and office of these muscles that a complete closure of the Glottis cannot be effected.

This disease generally arises from improper exposure to Cold or damp Atmosphere. Those persons who are exposed to great vicissitudes of the

weather are much more liable to the disease, than they who reside in Countries uniformly Cold - and it is on this account that Cases of Bronchitis are most generally met with in the United States during the winter & Spring. The slightest causes will produce this disease in those Children who have once had an attack of it - in such constitutions a common Catarrh will almost invariably be followed by the Croup - It is however a fortunate circumstance that second attacks of the disease are seldom so violent as the first. They nevertheless require the utmost caution on the part of the Physician.

Bronchitis has been ~~thought~~ thought by some very eminent Practitioners to be a contagious disease even the acute Dr Gregory seems inclined to adopt this opinion. There are however so far as I have been able to learn no evidences of the fact - there is on the contrary every reason to believe that the disease is not contagious. The same cause which produced it in one child of a family may give rise

to it, in another; and if we admit the contagious character of Trachitis we must extend the doctrine so as to include all those diseases consequent upon the application of cold to the Body. But where can that individual be found who will assert that Catarrh, Pneumonia or Rheumatism are contagious diseases? The same cause evidently produces all these diseases, and no reason can be assigned why any one of them is contagious and the other not. But whilst I deny the contagious character of this disease, I am willing to admit, that it may be founded on Hereditary Predisposition. Dr Potter asserts that it is as certainly an Hereditary disease, as Gout or Phthisis. I shall not attempt to discuss this question as it is one which can only be decided by experience, but shall proceed to enumerate some of the most common symptoms of the disease.

Some days previous to to an attack of the disease the child

appear inactive and fretful, the eyes are dull and heavy - symptoms of Catarrh succeed, a hoarseness attendes with a peculiar shrill and ringing sound in speaking and coughing, which has been happily compared to the barking of a small dog or Foxes. has also been supposed to resemble the noise from a Brazen Tube: Chillness sometimes attends the first stage of the disease, a pain is felt about the Larynx, great difficulty of breathing, accompanied with a whizzing sound in inspiration, the cough is commonly dry but if anything is spit up it is generally a purulent matter; sometimes however films are discharged resembling portions of a membrane together with these symptoms the pulse is hard and frequent the patient is restless and complains of an extreme uneasy sense of heat. The External Fauces on examination sometimes exhibit no appearance of Inflammation

most frequently, however a redness and even swelling is observable. It is remarkable that the natural functions as well as those of the Brain are seldom much disturbed. Children are often seen running about taking their food, indulging in the amusements peculiar to their age, apparently in good health whilst in reality the disease is rapidly advancing. if its progress be not promptly checked all the symptoms soon become much aggravated the inspiration is extremely laborious, the cough ~~difficult and~~ is troublesome and expectoration more difficult and in a few hours suffocation closes the scene.

The duration of the disease depends generally upon its violence and the means which have been employed for its cure; when no medical treatment has been resorted to, it almost frequently terminates fatally in about 24 or 36 hours

It is all important, that the alarming symptoms should be moderated during the first twelve hours of the disease; unless this can be effected the patient generally dies soon after. If however by the efforts of nature or by the assistance of art the child recovers from what has been called the second stage of the disease, the convalescence is always tedious, and attended with exfoliation of portions of membrane. In milder cases where the breathing is less difficult in the commencement, about the second day the skin becomes moist, the fever abates, the cough is less troublesome, and the voice gradually recovers its natural tone.

A peculiar morbid appearance is observable in the examination of those who have died of this disease. a membrane or tube of coagulable lymph, is found lining the whole extent of the Trachea descending even to the bifurcation of the Bronchia. Dr Potter says this body is found not only through out the Bronchial ramifications

but throughout many of their extreme ramifications into the smallest circuits as was beautifully illustrated in a subject prepared by the Dr for the demonstration of the late Professor Mitchell of Philadelphia. This continues the Dr is a secretion from the inflamed vessels of the part, and is of all consistencies from the smallest change from health to the secreting process perfectly formed in inflammation. The secretion of the more compact form to which the appellation of membrane has been given, seems to be the fibrine of the Blood semi-organized. Although this membrane is generally found in the Trachea of those who have died of Croup, still it is not to be considered either as a constant or necessary part of the disease. Dr Gregory has observed that its formation is always indicated by the manner in which the child breathes, throwing the head back so as to put the Trachea on the stretch.

With regard to the treatment of this disease there seems to be some difference of opinion

of opinion among Medical men. A
Pathological distinction has been made by
some, which is of no practical impor
tance. If there be such a thing at all
as a primary Spasmodic Croup, it must
invariably depend upon an Inflamma
tion of the muscles of the Glottis, and
therefore the same remedies are to
be employed in the first stage of
the disease, whether it be an inflam
matory or Spasmodic affection.

The Croup has been divided
into two stages; the first being that
of high inflammatory action. The
second is distinguished by the forma
tion of that preternatural memb
rane already spoken of. In the
first stage, it should be our ob
ject to subdue Inflammation
by all the means in our power and
thereby prevent the formation of that
membrane which "block up the pas
sage of air into the Lungs, inducing
suffocation and consequently death.
To accomplish this first indication
the most powerful antiphlogistic
remedies should be promptly em-

employed - our chief reliance is to be placed in general and Local Bleeding emetics, the warm bath, and purgatives If however we do not succeed in relieving the first stage of the disease, it then becomes our duty to endeavour to promote expectoration, and to support the strength of the system, which is generally much weakened by the previous depletory measures - for the purpose of answering these indications recourse may be had to to squills, Camphor opium, and other tonic medicines an occasional emetic may also be exhibited, and as a last resort Bronchotomy - has been recommended

I shall now say a word or two with regard to the particular remedies which have been recommended for the cure of this disease, and first of Bloodletting - This in my humble opinion is our Sheet Anchor It is never injurious: it may however sometimes be unnecessary In the milder form of the disease where there is not much inflam-

natory action, an Emetic will often supersede the necessity for the employment of any other remedy - One bleeding in the first stage of the disease is of more importance than half a dozen afterwards -

Emetics are always necessary and should never be omitted - for this purpose the Tartar Emetic or Ipecacuanha may be employed - If however after being given in very large doses they fail to produce the desired effect, the Corrosive Sublimate may be resorted to with the greatest advantage - The exhibiting of small doses or repeated doses of Calomel from the commencement of the disease, has been strongly recommended - "This practice says one in whose skill and experience I have an almost unlimited confidence" is rather calculated to injure the reputation of the medicine, than to cure the patient - It should be given in doses so large as to shock most of the

mid: Practitioners of the present day who are deterred from employing this valuable remedy to any useful extent by the fear of salivating their Patients Dr Potter says there is but little danger of producing a severe Ptyalism by Calomel in this disease. he has seen but one case out of several hundred. No one of these remedies should be relied on in violent cases to the exclusion of the others. Bloodletting being premised a judicious combination of Calomel & Tartar Emetic will generally cure the Patient. nor should the assistance of the warm bath be dispised. The application of Blisters to the Throat has been recommended. its propriety is I think very questionable. It seems to me that the irritation of the Blisters, may sometimes extend to the inflamed membrane, and consequently aggravate the disease. The decoction of Polygala Seneca has been lately introduced into the practice as a remedy for this disease by Dr Archer of Maryland. It is my opinion too stimulating to be of any advantage in

first stage of the disease - when however the inflammatory symptoms have ~~been~~ in a great degree ~~subsided~~ ~~etc~~ employment will ~~in my opinion~~ generally be attended with the most beneficial results - It is unnecessary for me to say anything with regard to the other medicines which have been recommended in this disease - in the first stage they ~~may be beneficial~~ must be injurious, in the last they may be useful - Bronchotomy is altogether inadmissible in Croup - for this simple reason, that we are unable thereby to remove the membrane which is the cause of the obstruction - and besides it prevents us from employing other remedies -

249

# An Inaugural Dissertation on Pulmonary Consumption.

Submitted to the Consideration of the Provost, Professors and Trustees in the University of Maryland for the degree of Doctor of Medicine by

Bennett Bussey of Maryland.

1828.

Pulmonary Consumption is a disease, concerning which a variety of opinions have been given, and even at the present day Physicians widely differ with regard to its Pathology; some contend that the disease never appears without the presence of tubercles in the lungs, while others say that Pulmonary Consumption can exist independent of a scrophulous predisposition.

As it would be not only tedious but also useless, to attempt an elaborate discussion of a subject, which probably may never be decided, I will merely submit my opinions with regard to the real nature of the disease, enumerate the principal symptoms and mention those remedies which appear to be most useful.

By the term "Pulmonary Consumption" I wish simply that disease, which is the consequence of Tubercles seated in the cellular connecting membrane of the lungs. Whether or not this term could be properly applied to other diseases of the lungs it is not for me to decide, but merely for the sake of precision; I am disposed to think, that the peculiar wasting of the body and of the lungs caused by the inflammation and suppuration of tubercles seated in the lungs, alone deserves the name of Pulmonary Consumption.

Those who have tubercles in the lungs are evidently of the strumous temperament, and it still remains to be proved

whether or not tubercles be ever formed in the lungs without an hereditary predisposition to them:
The strumous temperament appears under two modifications, the one is found in those who have, naturally, pale skins, loose flabby fibres and a sluggish pulse; the other in those who have ruddy complexions, firmer fibres and a brisk circulation. And in both modifications there is an unusual vivacability of the capillary arteries;
In children we can generally determine whether or not they are predisposed to Phthisis, by the smooth and rosy appearance of their cheeks, their fair hair and delicate blue eyes: the forehead is generally prominent, and it is remarked that the lips of scrophulous children are tumid;
As the child advances in years we perceive the swelling of different glands of the body, which as the patient progresses in age, progress in growth: and I am disposed to think that almost all of those diseases, which we hear spoken of as affecting different joints, are no more nor less than the effects of Scrophula;
As the general characters of Scrophula are so well known I will not longer dwell upon them, but proceed to describe its appearance in the form of Pulmonary Consumption

This disease seldom appears before the age of puberty, It is usually excited by causes which produce Inflammation. The patient complains of a weight or uneasiness in the chest, this sensation is attended by a cough which is generally dry, short and tickling, there appears to be considerable excitement, and the faces usually flushed, The pulse is frequent and sometimes tense.

As the disease progresses, the pain in the chest becomes more acute and often extends to one shoulder.

The cough becomes more troublesome, and at length the expectoration comes on: The lips of the patient are generally preternaturally florid.

At this stage of the disease hectic symptoms are more apparant, The patient is harrassed with evening fever and night sweats, These symptoms continue till the pus formed by the suppuration of the tubercles in the lungs has been expectorated, which in the early part of life will be accomplished in a short time.

The patient is then considerably relieved for that period and has few unpleasant or alarming symptoms until the same process of Inflammation and suppuration comes on. The patient, however is not entirely relieved from hectic symptoms; some cough and fever still remain.

As the patient still advances in life, the general health becomes more impaired, from the repeated struggles which he has survived, and his whole system becomes reduced from long continued hectic fever; nights are spent without repose, and each returning day presenting the same or a worse condition of his system, renders life almost a burden;

In the forming stage of Pulmonary Consumption, we have many remedies which exert a very beneficial influence in warding off the disease, but when Phthisis becomes confirmed very little can be accomplished by the Physician. In the forming stage or before inflammation of the tubercles has progressed in any considerable degree, Bloodletting is a remedy of inestimable value, Digitalis has been used with the same view but it has been observed "That the injurious effects of Digitalis often precede its good effects" and I would suppose that in every case of Consumption the lancet will accomplish every object which could be expected from Digitalis: Blood should be drawn in small quantities, and the operation should be frequently repeated, There are many cases of Phthisis in which mercury given with precaution, exerts the most salutary effects; cases in the forming stages, which are not attended with considerable action, have often been prevented from terminating in suppuration, by using

the Blue pill so as slightly to effect the gums.
Those who are predisposed to Pulmonary Consumption, should be very cautious in their dress. Flannel should be worn next the skin in winter by all, and in summer also, by the more delicate. It preserves the equilibrium of temperature, and prevents the system from being injured by every change of weather, and as our climate is so very precarious, it is my opinion, that this prescription should by no means be neglected.
Another important prescription, is Exercise; The Consumptive patient, in clear weather, should take exercise freely, particularly in the mornings, it gives tone to the digestive organs, and invigorates the whole system.
Attention should be paid to regimen and diet. The patient should take animal food in moderation, and it should be the principal article of food. All irritating and indigestible articles should be avoided.
Blisters or setons are almost indispensible in the prevention of this disease. A discharge from the surface should constantly be promoted by either a blister or a seton.
The cold Bath has also been highly recommended by some Physicians. But the most effectual means for preventing this disease is a Removal to a warm climate, and to a climate which is free

from many vicissitudes of weather.

These are the principal means which are used for the prevention of this disease. When Consumption becomes confirmed, as I before observed, very little can be done by the Physician. However we have some medicines which afford some relief. Opium is generally freely used in the latter stages of Consumption. When the patient passes nights without sleep, has great irritability, and is harassed by a continual cough, I am of opinion that Morphea might be given with great advantage as it possesses all the sedative but none of the stimulating qualities of opium. This also will answer every purpose for which Prussic acid has been given.

When the colliquative sweats are copious, the balsam Copaiba given as a Diuretic generally lessens the sweat.

As I intended merely to make some general remarks on this disease, I will end, after mentioning some diseases which may be mistaken for Pulmonary Consumption, and mentioning some differences which distinguish them from true Phthisis.—

Ulcerations in the trachea has often been mistaken for Pulmonary Consumption. This often arises from an insidious inflammation. It is attended with a tickling cough, the voice is slightly

changed, and uneasiness is experienced in some part of the Trachea; The breathing is sometimes oppressive, and there is generally slow increasing fever. Inflamation of the adjacent parts also very often exists at the same time; In the first stage of this affection mucus and lymph are expectorated, but as soon as ulceration takes place the sputa are mixed with pus; After this disease may have continued for some length of time fever is manifested but it is not attended with the copious night sweats which characterize the True Hectic. When Inflamation of the Trachea becomes chronic soreness is felt by pressing on the Trachea which sensation is not experienced in Pulmonary Consumption.

Chronic Inflamation of the Bronchia is another disease which may be mistaken for Consumption, It at first resembles a common Catarrh, but by the increasing fever and duration of the disease we are made sensible of its nature. When it has progressed so far as to make an abscess in the lungs themselves it is very difficult to distinguish it from Tubercular Consumption, In the first stage it may be distinguished by the wheezing and Catarrhal character of the disease and by the patient not wasting flesh and strength as always happens in Tubercular Consumption; The face

from the onset assumes a sickly pallidity, and the lips are of a leaden hue, whereas in Tubercular Consumption the face is alternately pale and flushed and the lips are beautifully tinged with a bright red.

When the more acute forms of Pleurisy have been subdued an obscure kind of increased action is often left which sometimes becomes chronic. This however may arise without being preceded by acute Inflamation. The patient has slow fever at night and has oppression in the chest. There is occasional cough which is most troublesome in the morning. Soreness will be felt in the integaments of the side effected, particularly if pressure be made. The disease progresses until dyspnea and emaciation are the two principal symptoms. In this stage it can be distinguished from Phthisis by the colour of the patients skin, and the difficulty of breathing which is increased by using the least exercise.

Chronic Inflamation of the lungs themselves has very often been mistaken for Tubercular Consumption. This disease in its progress bears a strong resemblance to Tubercular Consumption but a history of the patients life preceding the attack, and by procuring a knowledge of the health of the patients predecessors, we can generally ascertain whether or not the latter disease exists.

267

# An Inaugeral Dissertation on "Hydrocephalus Internus."

Presented for the consideration of the Faculty of Physic of the University of Maryland by

Charles F. Hughes of Baltimore

March 16th 1828.

To Alexander Glendinen M.D.

The following Inaugural Dissertation is most respectfully dedicated, as a tribute of sincere gratitude and high respect for the many kindnesses and valuable advice received from him by me, while pursuing my medical studies under his able guidance and fostering care

Chas. F. Hughes

March 16th 1828.

## Introduction

The many difficulties which present themselves to the mind of the writer, in selecting the subject of his Inaugural Essay, influence in a great degree his choice. It cannot however be expected that one who has neither age nor experience to support him, can produced anything new in relation to disease. — I shall therefore content myself by giving, as far as my limited abilities and time will enable me, a history of the causes, symptoms and treatment of the disease under consideration, known by the term Hydrocephalus Internus. Having chosen this subject for the following pages, it is my intention "safely to tread the beaten track;" and with this view shall quote the opinions advanced by Whytt, Quin, Rush & others.

# Hydrocephalus Internus.

The earliest correct description of this very interesting and too frequently fatal disease was, as well as I am informed given to the world by Whytt, whose observations on it, were not published until the year 1768; This valuable author describes with great accuracy the cause and advancement of the disease. —

Cullen in the edition of his nosology, published in 1785 recognizes this disease under the term Apoplexia Hydrocephalica.

Quin of Dublin, in his Inaugural Dissertation published at Edinburgh in 1779, and a subsequent Essay in 1790 describes Apoplexia Hydrocephalica with much more accuracy and proposes ideas altogether new as to its theory. —

Rush, our celebrated countryman not long afterwards adopted a similar theory of the disease to that of Quin, & in his works, has given us a paper in which he mentains the inflammatory nature of Hydrocephalus Internus. —

Whytt in speaking of its proximate cause, seems to make it depend upon a loss of balance between exhalation and absorption; And indeed this was the opinion of the

ancient as well as the modern writers, concerning Dropsies in other parts of the body.

Quin suggested the inflammatory nature of this disease in his Inaugural Essay written in 1779, and in a treatise subsequently, confirmed this view of the disease, by additional observations and numerous cases of dissection.

The same opinion was adopted by Rush, (med. Inq. & obs. Vol 2 page 193) about the same time, to which increasing weight has been given by discoveries on dissection, shewing the fortunate issue of many cases treated by Anti-phlogistic remedies.

The writer would venture to defend the opinions of the two last named authors; and for this purpose he conceives it will be necessary attentively to consider the causes of this disease, the symptoms and the mode of practice which has been found most successful in treating it.

Hydrocephalus Internus, may probably be divided into primary and secondary.

Rush enumerates the following causes "falls or bruises upon the head, certain positions of the body and childish plays, which tend to induce congestion, inflammation and subsequently an effusion of water in the brain." — Now we have every reason to conclude from these very valuable remarks, that falls, bruises or blows on the head, do produce inflammation

not only of the external integuments, but also the lining membrane of the cranium; which too frequently terminate in suppuration and gangrene. The other direct causes mentioned by Rush to which more might be added, especially the influence of the sun & of a reduced temperature, can scarcely be supposed to act in any other way than by increasing the quantity of blood in the vessels of the brain, and resulting in an inflammatory action in them. — The causes enumerated by Rush as acting indirectly in producing the disease under consideration, are surely fitted to produce an inflammatory ~~action~~ disease; these are states, either of inflammation or irritation in other parts of the system; and when the disease is thus produced, there is a mere translation of it, and not a change of morbid action; Surely, the membranes and the structure of the brain are different from that of other parts of the system, but then, they have not the power of producing an action in the vessels, sui generis. The effect of their operation therefore upon the brain must be an excitement of irritation & inflammation. These causes are, 1st Intermittents Remittent and continued fevers, 2nd Rheumatism, 3 Phthisis Pulmonalis, 4 Eruptive fevers & Worms. —

Having briefly noticed the causes of this disease, I propose next to take notice of the symptoms which attend it, and also of the appearances upon dissection after death, being further

proofs of its inflammatory nature.

The symptoms of Hydrocephalus Internus as related by Whytt appear to me to be as minute, at least, and as correct as those detailed by those who wrote subsequent to him, however, as the work of Quin is more modern and sufficiently correct, I will insert his description of the symptoms.

In general the patient is at first languid and inactive often drowsy and feverish, but at intervals, cheerful and apparently lively and free from any complaint. The appetite is bad, often nausea and vomiting occurs throughout the day, the skin hot and dry towards evening. Soon after these symptoms appear, the patient is attacked with sharp headache, chiefly in the forepart, and if not there, in the crown of the head, it is sometimes however confined to the side or sides of the heads, and in that case, when the posture of the body is erect, the head often inclines to one side, particularly that side affected. This headache is also frequently accompanied with an affection of the stomach, the vomiting being more troublesome when the pain is less violent and vice versa.

"The light is disagreeable to the patient at this period cries much, sleeps but little, and when he does sleep, grinds his teeth, picks his nose, appears to be uneasy, often screaming as if greatly terrified. The bowels in a majority of cases

are much confined, though it sometimes happens, that they are in an opposite state. The pulse in this early stage does not indicate any material derangement.

When the symptoms above mentioned have continued for a few days, subject as they always are in this disease to great fluctuation; the axis of one eye is generally found turned inwards to the nose, and the pupil upon that side rather more dilated than the other. Once when both eyes have their axes turned inwards, which sometimes happens, both pupils are observed to be larger than in health. The vomiting becomes more constant, and the headache more excruciating; at this period every symptom of fever makes its appearance; the pulse is frequent, the breathing very quick and exacerbations of fever take place towards evening, the face occasionally flushes, usually one cheek more than the other, temporary perspiration breaks out, but does not alleviate the condition of the patient. Epistaxis takes place about this period; also delirium and that of the most violent kind, particularly if the patient has arrived at the age of puberty, which together with all the preceding symptoms of fever, continues for a while to increase, until about the fourteenth day, often a much shorter space of time shall have elapsed since the appearance of the symptoms which were first mentioned in the above detail —

the disease undergoes that change which suddenly points out the commencement of what has been termed the second stage. The pulse becomes slow, but unequal, both as to its strength and the interval between the pulsations. The pain of the head or whatever part of the body had previously been affected, seems to abate, or at least the patient is less sensible to it, the interrupted slumbers or perpetual restlessness which prevailed during the earlier periods of the disease are succeeded by an almost lethargic torpor, the strabismus and dilitation of of the pupil increases, the patient lies with one or both eyes half closed, which when minutely examined will be found almost completely insensible to light, the vomiting ceases, and whatever food or medicine is offered, is usually swallowed with apparent voracity, the bowels at this period generally remain obstinately costive. ——

"If every effort made by art fail to raise the sinking powers of life, the symptoms of what has been called the 2 stage are soon succeeded by others which more certainly announce the approach of death. The pulse again becomes equal, but so weak and quick, that it is almost impossible to enumerate it. A difficulty of breathing nearly resembling stertor apoplecticus is often observed, sometimes the eyes are suffused with blood, the flushing of the face is more frequent than before, but of shorter duration, and followed

by a deadly paleness; red blotches sometimes appear on the body or limbs, deglutition becomes difficult and convulsions generally close the scene."

The symptoms no less than the causes of Hydrocephalus Internus, demonstrate its inflammatory nature. The intolerance of light the morbid sensibility of the auditory nerve, the delirium and flushings of the face can be explained upon no other principle. ——

One or two more points shall be noticed under this head in favour of the inflammatory nature of this disease; one first of the discoveries after dissection. — In resorting to the evidences of morbid action shewn by human dissection, especially when connected with a knowledge of the progress of the symptoms during the continuance of the disease, we may be considered as appealing to the highest source of information, with regard to the phenomena of diseases. Three particulars are to be noticed in attending to the appearances after death, — 1st, The thickning of the membranes of the brain the existance of inflammatory crusts, and the engorged & distended state of the blood vessels. The frequent occurrence of these things as consequences of inflammatory diseases of the Thorax and Abdomen, cannot or ought not to leave the least shadow of doubt with a candid and intelligent mind, that the same appearances of those decays of

Hydrocephalus arise from the same cause.

In speaking of the appearances on dissection of those who die of Phrenitis, a highly respectable author says "There is often a quantity of water effused into the ventricles." When we compare this appearance with those who have died of Hydrocephalus, we cannot help, I think, regarding this disease as a species of Phrenitis." (Wilson on febrile diseases 2d page 96) The effusion into the joints which follows acute Rheumatism, & those which succeed to inflammation of the Pleura, Peritoneum and other inflamed internal surfaces, would seem at least to give countenance to the supposition that the fluid in Hydrocephalus is produced by the same cause. — I admit that the analogy here presented does not unanswerably prove this to be the case, for by assuming this position we must deny that Dropsical affections ever do arise from any other cause than inflammation, which I am not prepared to do. — The thought just hinted at leads to the consideration of the last particular in relation to the appearances upon dissection; (3d) which is the peculiarity of the fluid, viz its refusal to coagulate by heat; and this property distinguishes it from ordinary Serum and this proves it to be a fluid thrown off from the circulating blood either by secretion or some equivalent process. Our Professor of the Theory and Practice has taught us that incoagulability is a property belonging to Dropsical effusions

into every cavity, and therefore infer that Dropsy, in whatsoever part of the body, it may occur, is an inflammatory disease. In Quin's treatise (appendix 2 case 12) there is an instance of fluid formed on the brain being coagulated by heat, With the exception of this single case, I neither know or ever heard of any circumstance to support the general doctrine.—

This property of incoagulability, as before mentioned distinguishes it very pointedly from serum. It's a charged fluid, and from the symptoms of the disease, of which it is an immediate consequence, we have sufficient reason to believe that it is the result of inflammatory action. This theory of the inflammatory nature of the disease under consideration, receives additional confirmation from the constitution and temperament, of the subject whom it most frequently attacks. It is, says Rathieger vol 1 page 71, a disease, that happens, so far as I have had an opportunity of observing, most commonly to lively, healthy, active children in whom of course acute diseases rather than of a low action, a debility may be supposed to prevail. The next point to be noticed according to the arrangement of this treatise, will be the means of cure, which have been found most effectual in the treatment of Hydrocephalus Internus. However, before entering upon this part of my subject, I will say a few words relative to a prevention of this fatal disease.—

The means of prevention may be classed under two general heads, each embracing many particulars, too numerous to be related here. The first general means of prevention is, avoiding all the direct causes which tend to excite the disease. Second; by endeavouring to remove as speedily and effectually as possible, all those diseases which act as marked causes, in producing it. — The first head of precaution, may be addressed to the nurse, or the patient, for this disease is not confined to children alone; and the second more properly is addressed to the Physician.

Having premised these reflections on the preventive method, I will now return to the consideration of the remedies which have been found most useful.

The first means to be noticed is bloodletting. And this exhibits another most decisive proof of the correctness of the theory adopted in this treatise. Venesection, though recognized by Quin in his list of remedies, was unfortunately but seldom resorted to by him, if we may be permitted to draw any conclusion from the cases, reported to have been treated by him. In seven cases successfully treated by our celebrated Rush, venesection was practised more or less in every case. This course should be adopted in the early part of the disease, and should be frequently repeated, always taking care to consult the pulse. Bloodletting from the

Scrofulous, pain has been recommended by some; this however, has its disadvantages. Blood drawn from the arm, in a proper manner, will reduce the general excitement as effectually and equally as quick, as it taken away from any other part. — Local bloodletting I conceive to be of great advantage — The great importance of this will be very evident, for while general bloodletting only reduces the force of the circulation, local bloodletting tends to relieve the distended and engorged state of the vessels of the part affected; which state is not only the cause of the affection so much to be dreaded, but also acts as the cause of excitement of the general system. — The second remedy is Blisters. — There has been and still is much diversity of opinion relative to the mode of operation of blisters among medical writers and practitioners. That they act as a stimulus to the part applied as well as generally, cannot be denied; I deem, tho' not the most important, is the abstraction of a portion of the more fluid part of the Blood. Rush says they are proper in every stage of Hydrocephalus Internus. In the early part of this disease their application should be preceded by general bleeding; and it is proper to keep up the discharge from the blistered surface for a short time, the better to secure your aim; and for this purpose the Ung.t Epispast. may be used to great advantage. Seton and Issues have been recommended by some; but in my

opinion they are too slow in their operation for a disease of this nature.

Cathartics, are the third class of remedies to be noticed. They should be assiduously employed from the commencement and the best agents for this purpose are, probably the cooling salts. They are more speedy in their operation and seem to occasion larger watery discharges. A combination of sulph. Pot. Potasse and pulv: Jalapa a Jalapa and nitrate of Potash will be found very useful. Calomel is more widely recommended by all writers. When used to procure a purgative effect, it should be combined with one or more of the above named remedies. In an advanced stage of the disease, mercurial friction may be resorted to, but should not be depended upon alone. As the bowels are disposed to be torpid in this disease, much larger doses will be requisite, than under ordinary circumstances. Cold applications come next to be considered. – Cold water, vinegar, and water, a pound of Ice put into a bladder, have all been found of much service in attracting heat from the head. The application of Ether, or other vaporizable fluids, by their sudden absorption of a large portion of caloric, are unquestionably calculated to produce a good effect. The position of the patient is by no means an unimportant

consideration. The head should be as much elevated as convenient; the room kept perfectly dark and very cool, much light being very injurious. All noise should be avoided in cases of morbid sensibility of hearing.

As premised in the introduction nothing original is pretended on my part. — In my humble opinion however the credit of first shewing this to be an inflammatory disease belongs of Quin. This Theory, plausible and defensive as it really is, would indeed merit but little estimation were it not that it leads to a system of practice, which experience has proved to be far more successful than any other which has been recommended. It is this correspondence between the Theory and the results of practice which gives to the former their only value.

By a course of strict investigation and close observation, Quin and Leech, have exhibited the true nature of the disease which has been the subject of this Essay, and they have instituted and recommended a course of practice which if attentively pursued promise more than any other to restrain the ravages of Hydrocephalus Internus.

303

305

An
Inaugural Dissertation
On
Hydrocephalus Internus
By
Edward H Lowndes

For
the Degree
Of Doctor of Medicine

Maryland. March 15th 1828

Perhaps there is no disease in Nosology which is described, by such a variety, of names, as the one which I have selected as the subject of my Inaugural Dissertation. Although we call Hydrocephalus a disease, it is more, from the tyranny of custom than strict pathological accuracy; as the derivation of the term implies, it is a collection of a watery fluid secreted in some part of the brain, and is the effect of a slow chronic inflamation of that organ. it is described by Doctor Cullen under the title of serous apoplexy, but the fluid which is found in the brain of those who have died of this affection does not resemble serum, except in appearance as it will not coagulate, by heat or any chemical agent; therefore we cannot

with strict propriety subscribe the neither
serous apoplexy; others again describe the
disease, as a chronic inflamation of the
brain, Renetis chronica vel subacuta, this
appellation is more suitable than any other
to the condition, in which the vessels are
in, at the time, the fluid is deposited
and is therefore, a more appropriate term
I think than any which has, as yet been
applied; Doctor Good in his excellent work
on the study of medicine; calls it Cephalitis
profunda, what his object could have been
for adding profunda, I cannot conceive
otherwise than he intended that the water
secretion always commenced in the
ventricles, which are situated in the interior
of the Brain; if that was the Doctors object
he is certainly correct, as that is the usual
situation of the incipient secretion.

although he may be correct as far as respects the seat of the disease; I cannot subscribe to his opinion, concerning the proximate cause the Doctor thinks debility to be, the proximate cause of this, as of all other forms of dropsy by debility I understand a relaxation of the exhalents, and consequently an infiltration of the serous or thinner parts of the blood through the patent mouths of the exhalents in this respect he does not differ from Doctor Cullen, except in the name of the affection, as both perfectly, as to the nature of the fluid agree; there are some who suppose the proximate cause of all dropsies to be a debility of the absorbents and among the advocates of this theory Doctor Darwin was the most prominent who is deservedly more celebrated for his poetical, than medical productions

It was at one time a popular opinion, but which is too often, but another name for popular error that this affection, was generally intimately associated with a scrophulous diathesis of the general system, there are instances of congenital Hydrocephalus, where a separation of the cranial bones has taken place to an almost incredible distance, and children under such circumstances, have lived in apparent ease and comfort for several years.

Such occurances as these are sometimes supposed to depend upon, an hereditary scrofulous taint but it is a subject as yet sub judice, and would afford abundant materials, for the theoretic speculations of those, whose genius may unravel the mysterious and almost inscrutable intricacies of its causes, such instances are comparatively rare, and generally

come under the cognisance of the obstetrical
the few remaining advocates of the impe[illegible] tradition
elies of the gastric pathology, think hydro
cephalus to be frequently connected with a
disordered state of the chilopoietic viscera
when this is the case, the Braen I imagine
s predisposed, and the congestion of the diseas
d rises, excites the action of the heart, and
the, Braen being the weakest part, it consequ
ntly must be the first to suffer, and when it
loes, the vessels take on a secretory action, and
leposite the fluid in the ventricles. the presence
of worms in the intestinal canal, occasion
ally give rise to symptoms of Hydrocephalu
but such symptoms are speedily removed, by
the exhibition of proper Anthelmintic
medicines. our illustrious countryman Doctor
Rush who is deservedly acknowledged, to have

written one of the best chapters, in his enquiries
on Hydrocephalus, that has ever been presented
to the medical world, he thinks, that in the
very highly acute form of Phrenitis, the inflammation
is so great, as to transcend the grade
of secretion, as is frequently the case in cynanche
trachealis, and hence the term cynanche sicca
the Doctor thinks that the vessels of the Brain
are in a condition, intermediate between that
of serous apoplexy, and the high inflammatory stage
of Phrenitis, consequently they are labouring under
chronic inflammation; a condition which is
certainly the most favourable for that secretory
process which takes place, at the time the
fluid is deposited; now if we impartially take
into consideration the causes which operate
on the brain in producing this disease, and the
nature of the fluid, which is deposited

we must certainly view Hydrocephalus to be the effect of a chronic inflammation, wherein the vessels assume the function of secreting vessels as they frequently do in chronic rheumatism producing permanent lameness; the greatest source of error in the treatment of this disease with many Practitioners was in considering it a disease of debility; but the great oppressent langour and debility, which attends those laboring under it, is always partially removed by bloodletting: equally as much blood can be frequently abstracted in such cases as in gastritis where the pulse is not the surest and most infallible criterion before bloodletting, we cannot for a moment suppose that it is debility which induces the patient to complain in a low muttering tone of his head, debility would not in the first instance

cause the pupil to contract, and the retina to be morbidly sensible to light; as the disease advances, and the pressure on the Brain increases, the origin of the nerves must be equally affected, and deprived of their usual energy and power, the pupil then dilates, and the retina is capable of bearing a stronger light. From the records of medicine, and personal observation, it would appear, that this disease was in some families constitutional; the vessels of the Brain in those persons so affected, I imagine are in a very excitable state, and the first exciting cause, which is applied induces them to take on a secreting action as this disease is always in my opinion produced by the usual causes of inflammation; and most frequently succeeds

those inflammatory fevers, which have not been properly treated by bloodletting, and other antiphlogistic remedies; where there is a great determination to the head. I am confident I do not stand alone, in considering chronic inflammation as exclusively its proximate cause; and in considering the disease as the effects of inflammation I shall adopt a course of treatment strictly antiphlogistic, as the conclusion of my thesis; which I think is applicable to it, according to the views I have taken, and the theory I have adopted of its cause.

Although the absorbents are never primarily affected, in the incipient stages of dropsy they are concerned in its cure and that actively from the circumstance of their taking up the fluid which is deposited, and

and removing it from the system by the different emunctories; and from the circumstance of the doubt, which has, for a long time agitated the votters of medical science, as to the presence of absorbents in the Brain, this formidable affection has been looked upon as an incurable disease, although, the diligent and persevering Anatomist has failed to prove the existence of lymphatics in the Brain by ocular demonstration; nevertheless can we for a moment doubt their existence we certainly admit the Brain, the supposed seat of all our intellectual operations, to be an organized body, and as such it must be endowed with lymphatics.

But we have a stronger proof than this in favour of our position; have we not both read and heard of large portions of Bra

being removed, and what other outlet could there have been, except it was through the medium of the Lymphatic System; then if large portions of disorganized Brain could be removed, and without the least detriment to the Patient, surely, the watery secretion can be, which very much disturbs the general health; our object, should then be in the first place, to lessen the action of the heart and thereby, put a check to this secretory action which is going on in the vessels of the Brain, which is most promptly and effectually done by the lancet, we should not be intimidated, ~~at~~ on viewing the depression which the patient, is apparently almost overcome with, such symptoms usually attend compression of the Brain from other causes

such as extravasated blood, and collections of matter within the encephalon; because we cannot draw ten or twelve ounces of blood from a child labouring under this disease, we should not, on that account refuse to draw any; how many valuable lives have been rescued from the confines of the grave, by the abstraction of four or five ounces of blood, by a judicious practitioner. If we turn our attention to the state of the circulation, we will generally find it to be in that condition, which imperiously demands the use of the lancet; we should not be deterred, by small pulses provided they be hard and tense, from drawing blood particularly in cephalic affections, as they almost invariably improve, after the operation has been performed; after bloodletting

has been carried to its proper extent, a ques-
tion naturally arises, have we no auxiliaries
to it, I would answer in the affirmative
calomel from its well known power of
promoting absorption, would be an excellent adju-
vant in the treatment of this formid-
ble affection, especially when it is carried
to that extent as to affect the salivary
glands; more have recovered from this disea
under a salivation, premised by proper evac-
uations by the lancet; than under any
other mode treatment; issues too, I imag-
ine might also assist in the cure of this
disease, but too much reliance, should not
be placed on them in such a formidable
affection

338

335

A Dissertation on
Pneumonia
Submitted to the
Professors and Trustees of the
University of Maryland
For the degree of Doctor of Physic
By Leander W Goldsborough
Of Frederick City Md.
March 15th 1828

To Dr Charles H Goldsborough
This essay is affectionately inscribed
as a testimonial of gratitude for
his care and attention whilst super:
intending the authors advancement
to professional life.

Introduction. The difficulty of writing without experience and the hazard of attempting subjects about which there exists great diversity of opinion is sufficient to make every one cautious in selecting the subject of his inaugural thesis. Influenced by these considerations we have fixed upon a disease which although pretty well understood is still possessed of sufficient interest to require the attentions of physicians. As to originality we claim none, none can be expected, all we wish to do and all we shall attempt is to treat the disease as it has been treated and to point out what we shall presume to consider as errors too generally received.

# Pneumonia.

Pneumonia derived from pneumon a lung is a genus of disease in the class pirexiæ and order phlegmasiæ of Dr Cullen. Under this title he comprehends "the whole of the infla:mations affecting either the viscera of the thorax or the membrane lining the interior surface of that cavity." This is certainly the most judi:cious plan of treating the matter, for had he attempted to distinguish between the varieties of different authors he must have failed, but if he had succeeded, what advantages would have resulted from it? In a prac:tical point of view none. The rapidity with inflamation of the serous membranes spread is itself sufficient to upset the

notion of giving names in this disease derived from the part supposed to be affected. As to the term Peripneumony athough some may doubt the existance of such a case, it is more than probable that it frequently occurs, and is always characterized by greater violence than the inflamation of the pleura. But as we cannot always distinguish the one from the other and as perhaps in neither case is the inflamation exclusively confined to either the substance of the lungs or their investing membrane the word might be abolished; or if retained, retained only to express a more violent pneumonia.

SYMPTOMS. This disease comes on with occasional chills and flushes of heat, accompa:
nied

with the usual disagreable sensations ushering in all febrile affections. The patient becomes restless, the pulse quick and the skin preter. naturally warm. He begins to cough and complains of pain in some part of the thorax

These symptoms continue to increase the skin becomes hot, the pulse quick strong hard and full. The cough which at first was moist is now dry and excessively painful

The breathing is difficult and irregular in consequence of the pain being increased by a full inspiration. Which (the pain) is now completely developed. It is not always fixed, sometimes it is felt in one part of the thorax and sometimes in another. It is generally more severe on one side, frequently the right

but as often on the left, in either case the patient cannot lie well on the side affected. As in all violent inflamations the secretion are ~~are~~ checked, the tongue is furred and the urine high-coloured. Having detailed the symptoms as well as I could we shall proceed to say a few words about the

Causes. This disease most frequently happens to persons of vigorous constitutions, and those who are much exposed to the vicissitudes of weather. It most frequently happens in the spring though it may occur at any season of the year and to all ages sexes and conditions. Cold applied to the body in some form or other is univer: ally admitted as the most frequent remote cause, though it may arise from others as excess

exertion of the lungs, drinking cold water when warm and perspiring is mentioned in Johnson's journel as having caused the disease.

Terminations. The most frequent termination of pneumonia is by resolution this may be known by the gradual manner in which the pulse is restored to its natural standard at the same time that the cough pain and difficulty of breathing subside with the return of the different secretions. Suppuration is another termination of this disease; the tendency to which may be ascertained by the pulse becoming softer and more frequent by the patient being affected with slight chilly sensations with a pale and anxious countenance. The pain is less acute and

when togather with these symptoms or rather as they advance, the difficulty of breathing continues the pain subsides or is remarkably dull and obscure with a weighty sensation in the side we may generally infer the existance of matter. Other terminations of pneumonia are spoken of as gangrene which is very rare.

A much more frequent when it ends fatally is effusion of blood and serum into the bronchia and cellular texture of the lungs producing suffocation.

Treatment. The first step in the treatment consists in copious bloodletting. This is the most powerful means we have of reducing the excitement equalizing the circulation and preparing the system for other remedies.

By drawing of a sufficiency of the circulating fluid we prevent many ill consequences ~~too~~ too

frequently the result of a protracted pneumonia.
Besides all these advantages it relieves imme:
:diately the violent pain which the patient
suffers, relaxes the surface of the body, promotes
expectoration and the operation of sweating
medicines, the advantages of which you
cannot always profit by unless preceeded
by bleeding. Some writers talk much about
the cautious use of the lancet in this disease,
they tell us that we should be governed by
the age constitution and condition of the
patient, that the quantity necessary to be
taken from one man would sink another
D.P. Now the fact of the business is that although
we admit in part the correctness of this reason:
ing we say that as a general rule the pulse
is the only criterion by which we are to be
governed; every body knows that an old man
will not bear the loss of as much blood as a young

ne, neither is as much required for the very reason that he does not so rapidly redeem it. So long as the pulse is hard and the pain and difficulty of breathing continues we must bleed.

The blood should be drawn from a large orifice in order that a dicided impresion be made at once. The physician should keep his fingers upon the pulse and not be satisfied untill it is reduced below the natural standard. If it be the first or second bleeding and the pulse very high and bounding he should carry it to the extent of inducing sickness or fainting, for in this disease as in all inflammatory cases the pulse will rise and become more irresistable unless you at once adopt decisive measures. Generally two such attacks with the lancet will be sufficient to bring the

ase under the control of milder measures and as
hneumonia is sometimes cured by expectorants and
diaphoretics we shall proceed to some remarks
about them. And first of Digitalis. This
article has been employed by Dr Barton and
others with a view to its expectorant and seda-
tive effects. We should certainly not expect
much advantage from it as an expectorant
it has not we believe answered the purpose
of physicians generally, besides the theory
of its operation is perhaps more obscure
than that of any other article of the class.
As to its power of controlling the action of
the arterial system I we know nothing, never
having it used the difficulty however of
egulating the dose and the admitted supe-
riority of other means are sufficient reasons
or abandoning its use. The best expectorant
ndoubtedly is antimony. This may be given

combination with nitre in the dose of a quarter of a grain of the former with eight or ten of the latter every two hours or so. Antimony is said by some to be beneficial by its sedatives qualities independant of its neauseating effects. That this articel acts directly as a sedative is a new idea, it may be correct, though we believe we have seen it injurious in high inflammatory fevers where its application had not been premised by bleeding. If in the advanced stage of the disease there is still pain with a difficulty of ~~breath~~ expectoration and a weak pulse, one or two grains of Camphor added to each of the above powders will be of great service, especially if aided by the action of a blister. After sufficiently reducing vascular action some gentlemen are in the habit of giving antimony alone in teped water so as to produce copious diaphoresis

if not to obtain some emetic effect. This is said
to be an excellent practice though perhaps alter-
nate doses of nitre would be an improvement.
In the low stage of pneumonia hinted at in
the preceding sentance Dr Potter states that there
a diminished susceptability in the nerves of
the lungs and bronchia and recommends mercury
though he says that squills, seneka and the balsams
are been given with great benefit. The Eupato-
rium Perfoliatum has proved highly serviceable
in this typhoid condition of the system. So
also is the Asclepias Tuberosa, given ~~in~~ as Dr
Thacher recommends in the form of a strong infusion
That physician states that it possesses "the peculi-
and almost specific quality of acting on the organs
of respiration, powerfully promoting the sup-
pressed expectoration and thereby relieving the
breathing of pleuritic patients in the most
advanced stage of the disease". Inhalations are
said to be good in pulmonic affections the

tam of hot water, vinegar and of the decoctions of Chamo:
le flowers are mostly used in pneumonia after the
olance of the inflammatory action has abated. While
e patient is under the use of these medacines he should
rink freely of flaxseed tea, barley water and solutions
Gum Arabac. In the last stage of the disease
hen the cough continues with but little pain
nd no fever, a mixture of Gum Arabac. Tinct Opii
nd liquorice in warm water is a very admirable
reperation. The temperature of the patients room
hould be carefully regulated. Dr Cullen says
t should never exceed 60 degrees of Fahrenheit's
hermometer. His diet must also be attended to. It
hould be of the lightest and most digestible
ind, His drinks should be asidulated and
oderately warm, and finally the whole of the
ntiphlogistic regimen ought to be rigirously
nforced. We now come to speak of blisters in

is disease. They are in a few cases unnecessary though
much oftener indispensable, particularly where the
patient is much weakened by morbid action before the
physician sees him, in which case it is not always
possible to draw a sufficiency of blood to supercede
their use. They should never be applied when there
is much action, but after two or three bleedings they
answer every indication. A fine large one should
be applied immediately over the seat of the pain
and suffered to remain on twelve hours. By
the proper use of blisters we frequently prevent
chronic coughs and other dangerous affections of
the thoracic viscera. We shall now conclude
with a few remarks on the use of Cathartics.
Almost all writers agree that medicines of this discription
will not answer in thoracic affections. The utility of keeping
the bowels regularly open with cooling laxatives is agreed
upon by all. Perhaps they might be carried to a
greater extent. We think we have seen purging
pretty fully induced in the first stage of the disease

th manifest advantage, the expectoration was very little any diminished. But the truth is, in this stage of e disease the expectoration is always ~~is always~~ earing whether you use purgatives or not. As we now of nothing to which we can refer this circumstance, ess it be excess of action, why not use this most powerful auxiliary to bloodletting untill we have sufficiently lowered the tone of the pulse to trust the ure to other means? However we are not very enacious upon the subject, we merely propose the uestion, with a hope that we shall some day or ther be better able to judge after a more extensive xamination of that best of all books EXPERIENCE.

369

To

Doctor W. H. Glendinen,

This Dissertation, is

respectfully inscribd;

By one, who,

Has had the Honor,

of being a pupil,

To so kind and gratefull; a

Preceptor.

March 20th 1828.—

# A Dissertation on the Atmosphere

Since the days of the alchemists it is well known that the science of Chemistry has been considered in a new light, and as subservient to very valuable and usefull purposes to mankind in general. The minds of the learned and scientific struck with the beauty and importance of its resources entered into the field of research with unmitigated ardour; and the present improved state of the science tell us what they realliz'd.

In the period between the years 1772. 1787 a new mine of experimental research which promised the most curious and interesting results had been opened at first by the genius of Dr. Black and allready pursued with much sagacity and industry by Dr. Priestly. But it was Lavoisier of France one of the most celebrated chemical philosophers of his time; a man whose views were far above any pecuniary consideration, who succeeded the former gentlemen in their particular views, and who devoted the last days of his life in experimental research that held the most importance on the branch of Chemistry; of which we are about

to make some brief remarks. We allude now to the discovery of the properties of certain aëriform substances, gases, or (as they have been called) factitious airs. They are bodies which Chemists have called elementary or bodies which are not in the present state of Chemical science considered as compound. They are bodies which have never been insulated they have been found only in combination with other bodies; and which had hitherto escaped the attention of Chemical inquirers.

After having said thus much as a desultory introduction to the after part of our dissertation; we will remark that it is our purpose to make some general observations on the atmosphere or the common air which surrounds our globe. As it would be very proper we think and for the better understanding of the subject we shall proceed analytically; and first with,

## Oxygen.

This elementary body forty or fifty years ago was unknown to the scientific world; it fortunately happened however to Dr. Priestly in the year 1774, when going through the experiment of heating the red powder which was called the precipitate per se, (but now more properly called by the name of the red oxide of mercury) in a glass vessel; a quantity of air was driven off from the powder which the mer-

-cury assumed its metallic form; he collected this air in a convenient vessel and examined it; he found it possessed of certain properties different from the atmosphere and in one respect he said it was much purer than the common air; he called it from the theory of the time dephlogisticated air.

Scheele a Swedish chemist shortly afterwards discovered this same gas without any previous knowledge of Priestly's discovery; he called it Empyreal air; it was also called Vital Air by some. Soon after this Dr. Priestly communicated his experiment to Mr. Lavoisier who with great ardour investigated the truth of Priestly's experiments; and from his investigations and observations are we indebted for the explanation of various phenomena: Lavoisier shewed that the mercurial precipitate per se, by being heated in a retort, and the retort being adapted to the pneumatic trough the beak of which terminated under a glass bottle filled with water; that an air passed over from the retort into the glass bottle, which was rendered evident by the displacement of the water in the vessel; and also a very important change had been effected in the red powder, which had previously been noticed by Priestly which was that it had resumed its metallic lustre and brilliancy and consequently was no longer a red powder but in fact running mercury. But the mon-

-der did not end here; the ingenuity and curiosity of the able Chemist led him to examine the air which he had collected: he did so, and found that it was an air highly respirable and supporting combustion in an eminent degree; for by putting a lighted candle or any other burning body in a jar of the air it burnt with increased brilliancy and force. He called it Oxygen gas: Oxygen in combination with the matter of heat or Calorics, for we do not speak of the gas in the abstract but in combination with a principle for which it has a great tendency to combine and to pass into a gaseous state.

This gas can procured in a variety of different ways and from a variety of substances.
1. From the black oxide of Manganese heated to ignition in an iron retort; or by mixing the same powder with twice its weight of Sulphuric acid in a glass retort and applying the heat of a lamp; very pure Oxygen gas can be procured in this manner.
2. It can be obtained from Saltpetre; by putting a quantity of this salt in a coated glass retort and applying a red heat Oxygen gas is formed. But there are two objections to this mode of preparation: for first the substance which remains in the retort can never without a great deal of trouble be got out,

ing another circumstance is if you are not very carefull in pushing the operation at the end to obtain the greatest portion of gas, another substance will come over so as to render the gas so produced very impure. A great quantity of the gas is generated in this way for it is said one ounce of salt yields 1300 cubic inches of gas.

3. It may also be procured from the Oxy-muriate of Potash a salt that yields it in great abundance; by heating it in a small glass retort over an argandts lamp. The gas obtained in this manner is much purer than that produced by any other process. These are the principle substances from which Oxygen gas is obtained: but it can be obtained in a variety of other ways which it is not necessary to mention at this time. It should be observed that very oxygen gas is generated by immersing some green vegetables (such as the leaves of mint) in a vessel of water covered and exposed to the rays of the sun; in a short time small bubbles will have collected over the surfaces of the leaves and subsequently rise through the water to the surface. —

The gas when produced by any of the above processes is inodorous, colourless and therefore transparent like the atmosphere; it is heavier than the atmosphere according to Sir H. Davy 100 cubic inches at 60° Fahrenheit weigh 34 grs.

All combustibles burn in oxygen gas with greatly

increased splendour or in other words it is eminently a supporter of combustion.
It was once supposed that Oxygen gas alone was necessary to combustion; but it has since been found out that other bodies beside Oxygen gas have an equal right to be considered as supporters of combustion.

Oxygen gas is very necessary for the support of animal life; without the use of this gas every breathing creature would cease to exist.

This gas when respired produces all the effects of a powerfull stimulant; it at first increases the strength and vigour of the body producing a degree of burning heat in the lungs; after continuing to respire it for a long time the stimulus of the gas is more than the animal body can bear; Vertigo, Coma, Convulsions and death eventually would be the consequence no doubt from the long continued respiration of this gas; yet it is a fact well known that a mouse, bird, or other other small animal will live four or five times longer in a vessel of Oxygen gas than in one filled with atmospheric air of the same dimension.

But a circumstance of great moment and one of the highest importance to us, is the relation which this gas bears to the function of respiration; for the perfect performance of which it seems necessary that Oxygen gas should be inhaled into the lungs there to effect

a very important change in the dark colour venous blood so as to render it fit for the usefulness of the animal economy.

It appears that Dr. Priestly was the first of the modern chemists who turned his attention to respiration and he concluded from his experiments that the blood as it passed through the lungs gave out phlogiston to the air which was expired loaded with that substance and of consequence that the purpose of respiration was to free the blood of phlogiston.

According to Lavoisier the blood absorbs no air in the lungs but it gives out hydrogen and carbon which combining with the oxygen of the air inspired form water and carbonic acid. Now there has been an objection to this hypothesis of Lavoisiers in as much as the quantity of carbonic acid formed is exactly equal to the bulk of the oxygen which disappears; therefore this oxygen must be changed into carbonic acid in the lungs for it is known that oxygen when changed into carbonic acid does not sensibly alter its bulk. From this it has been concluded by Dr. Thompson that the blood must emit carbon and that to an extent of about 3/4ths. of a pound in a day: this he considers is all that happens in the lungs: the watery vapour he allows is secreted from the blood to mix with the air expired;

but probably the secretion takes place in appropri-
-ate organs. Then by the loss of Carbon Venous
is changed into arterial blood.

Our distinguished professor of Chemistry Dr.
De Butts with his usual ingenuity accounts
for the phenomena in a different way.
From the doctor's view of the function of di-
-gestion &c; he concludes that the colour
of venous blood is owing to a portion of the
Carburet of iron and as it has been proved
that the colour of arterial blood is depen-
-dependent on a portion of the phosphate
of iron it is reasonable to suppose that the
object of respiration is to change venous
into arterial blood for that purpose is
the blood carried to the lungs from the right
-side of the heart loaded with carburet of
iron and also having in connection with it
a quantity of free phosphoric acid: when this
blood then arrives in the lungs it is exposed to the
contact and action of atmospheric oxygen the
moment the oxygen comes in contact with the
venous blood it combines with the carbon which
is expired in the state of carbonic acid; the
liberated air then finding free phosphoric
acid (which is always found in venous blood)
combines with it and in that state passes with
the other ingredients constituting arterial
blood into the left side of the heart.

Oxygen in combination with the metals form
a class of substances called Oxides. Carbon,
Sulphur, Phosphorous &c combine with ox-

-ygen so as to form a very interesting class of substan-
-ces called acids; some of the metals are capable of
combining with a very great proportion of Oxygen
and thus forming not only oxides but acids.
From these circumstances the French Chemists
considered Oxygen as the only acidifying prin-
-ciple this is not the fact for in the progress
of the science it has been found that there
are other principles that are the cause of acid-
-ity in some cases without the interference of
Oxygen at all.

Oxygen is an essential constituent of atmos-
-pheric air, of water, of saline and earthy bodies;
it is also a principle of animal and vegeta-
-ble matter. From all this it would ap-
-pear that the relations of this principle are
more numerous than any other body known.-

## Nitrogen.

This gas had ~~been~~ been considered as an
elementary gaseous body until the time
of Berzelius who with many experiments
and theoretical reasonings endeavoured to
prove that it was a compound body com-
-posed of Oxygen and an unknown base
for which he proposed the name of Nitri-
-cum. This base however is considered
by some and perhaps with great proprie-
-ty as purely hypothetical and as it never

has been produced in an insulated state Chemists cannot at present know any thing of its properties. A series of experiments to prove that nitrogen was a compound body had been got up by Mr. Siers of London who precisely agreed with Berzelius in his views of the subject. Sir H. Davy's attention was next directed to the subject on the presumption that nitrogen was an oxide; which however was not attended with any better success than those of the former gentlemen. The general statement of the inquirers therefore have been found to lend no strength to the supposition that nitrogen like oxygen is any more than a simple elementary body according to the present state of chemical knowledge.

Nitrogen is said to have been discovered by Dr. Rutherford in the year 1772. It was found that after seperating all the oxygen of any quantity of the atmosphere that there remained a gas which possessed peculiar properties and differed from oxygen gas or any other kind of gas which had hitherto been discovered. It was discovered that when a small animal was inclosed in a glass vessel containing common atmospheric air and the vessel being stopped tight; that after a certain period the animal would die, and inverting the mouth of the vessel into water and then taking out the stopper; that the water would rise and occupy a certain portion of the vessel: the air that remained was examined; and, it was found to be a gas

unfit for respiration or combustion; it was also found that fixed air was formed in this case. Now it is not to be supposed that all the air which was left was fixed air on the contrary there was comparatively but a very small quantity; for on separating the whole of the fixed air by lime-water it was found that just 1/5th. of the certain portion of air had disappeared and that the remainder 4/5ths. was that air which we have just mentioned as unfit for respiration or combustion: it was called azotic gas, azote; a name given it by Lavoisier from the circumstance of its not supporting animal life.

Now it is well known that there are various gases which are unfit for the support of animal life beside this one; for this reason the name Azote was dropped and the term Nitrogen has been substituted from the circumstance of its having been found to exist in a number of nitrous preparations.

Nitrogen gas may be procured though not absolutely pure yet sufficiently so for the purpose of exhibiting its general properties in either of the following manners.

1. Mix equal weights of iron filings and sulphur into a paste with water and place the mixture in proper vessel over water supported on a stand; then invert over it a jar full of common air and allow this to stand exposed to the mixture for a

day or two. The air contained in the jar will gradually diminish as will appear from the ascent of the water within the jar till at last only about 4/5 ths. of its original bulk will remain. The vessel containing the iron and sulphur must next be removed by withdrawing it through the water, and the remaining air may be made the subject of experiment. 2. By burning phosphorous in a glass jar containing common air the phosphorous will combine with the oxygen and form phosphoric acid which will fill the jar with dense white fumes, after a short time these fumes will be condensed and the remaining air will be tolerably pure nitrogen gas. 3. Nitrogen gas can also be procured from the lean part of flesh meat; put into a glass retort the lean of beef cut into small pieces and pour on it very dilute nitric acid then connect the retort with the pneumatic tub and apply a heat of 100°. gas will be disengaged and is collected in the gas bottle. This gas as well as oxygen is not absorbed or not in any very appreciable quantity by water. It is lighter than the atmosphere 100 cubic inches according to Sir H. Davy at 60° Farenheit weigh 29.37 grains, consequently it is also lighter than oxygen gas. It is not a supporter of animal life; for, it immediately proves fatal to animals that are confined in it. It is also a non supporter of combustion for it immediately extinguishes all burning bodies when they are

immersed in a gas containing it.
Nitrogen is susceptible of combination with various bodies and the compounds possess in many instances remarkable properties.

In the proportions of four parts of nitrogen with one of oxygen it composes a mixture resembling the atmosphere in all its properties: for it can be respired with safety for any length of time by an animal and burning bodies burn in it as in the atmosphere.

In combination with different proportions of oxygen it forms nitrous and nitric acids. It is also essential to animal matter; from the putrefactive decomposition of animal matters nitrogen and another elementary principle hydrogen are evolved; they combine and form the Volatile Alkali or ammonia. Nitrogen in combination with Carbon forms a carburet of nitrogen or Cyanogen which compound is the base of the Hydro Cyanic acid (Cyanogen in combination with hydrogen) or prussic acid as it is still called which is well known as one of the most virulent of the vegetable poisons. Although nitrogen is a necessary constituent of animal matter yet it appears to form a very small ingredient indeed as an ultimate principle of vegetable matter. From this it appears that nitrogen bears great relations to other

bodies and is little inferior to oxygen in point of importance in the great theatre of scientific pursuit.

## The Atmosphere.

It appears then from the facts stated in the preceding pages that the atmosphere is a compound or mixture of two different gases.
It is to Dr. Scheele that we are indebted for the fact that the atmosphere is a compound: for it is well known that to his time it was considered as an elementary body: Fire, air, earth and water being considered as four elements. Yet it has come to pass even before the present day that those bodies that ~~then~~ were once considered as elementary have been ranked as compound bodies and on the contrary those that were once considered as compound bodies are now placed among the simple elementary bodies.

This celebrated and illustrious Chemist Dr. Scheele in the course of some experiments after he had found out oxygen gas had reason to infer that the atmosphere was not a simple body. He took a glass bottle open with the bottom from it and he placed it on the shelf of the pneumatic tub and in this glass vessel he placed a small quantity of

of the substance called sulphuret of potash; it was supported on a stand above the surface of the water of the trough in the Vessel. After letting it remain in this situation for a time he found that a portion of the air in the vessel was taken away & for the water in the pneumatic tube had raised to a certain height in the Vessel. He after this took out the sulphuret of potash and examined it he found that it was no longer a sulphuret of potash but a sulphate of potash: potash containing sulphuric acid. Dr. Scheele found in this experiment that just 1/5th. of the air which was originally contained in the Vessel had disappeared and the remainder was no longer fit for respiration or Combustion.

The analysis of the atmosphere was most satisfactorily demonstrated by Lavoisier by the following experiment as explained in his Elements of Chemistry more enlarged. He took a glass vessel shaped like a matrass placed in it a small quantity of quicksilver and connected the superior part of the vessel with a right angular tube the one end of which being connected with the matrass and the other end with a gas bottle which was placed on the shelf of the pneumatic tub

He took care that the whole apparatus should be air tight and rule adjusted in its place.

He then had a portion of mercury in the matrass and all the parts ~~occupied~~ of the apparatus except that occupied by the small portion of mercury was necessarily filled with the atmosphere; an argands lamp was placed under the vessel containing the mercury and after keeping up a gentle heat for about 12 days he found that the mercury had suffered a considerable change, a part of it was no longer a shining fluid metal but was changed into a red powder; and also the water of the pneumatic tub had rose in the gas bottle and occupied a certain height in it. Lavoisier then examined the air which remained in the apparatus after the experiment and found that it was unfit for respiration or combustion. He next collected the red powder which was in the matrass and distilled it in a retort by which operation the mercury was regenerated it appeared in fact in its original metallic state and an air was given out which being collected in a gas bottle and examined he found that it indicated the usual phenomena of supporting combustion respiration, &c; and that it was the same gas which Dr. Priestly had discovered, viz; oxygen gas.

These results then afford the

most satisfactory evidence that the atmosphere is composed of two distinct fluids. The one is capable of giving its base to mercury, and when separate is eminently adapted to the support of animal life and combustion; the other has no affinity for mercury and does not possess the other important qualities. The object then of Lavoisier was to calculate the proportions which these two gases bore to each other; it was very easy to measure them and accordingly he did so. His report was that the atmosphere was composed of 1/4th. oxygen gas and the other 3/4 ths. of mephitic air. — But it appears that Lavoisier was not altogether correct in his calculations for according to the experiments of the best Chemists the atmosphere is now found to be composed of 1/5 th. of oxygen gas or 21 parts in the hundred and the remainder 4/5 ths. or 79 parts in the hundred is azote or nitrogen gas as it is more properly called.

It would seem then that oxygen gas is the only ingredient on which the chemical effects of the atmosphere depend. Hence combustible bodies burn in the atmosphere only in consequence of the oxygen it contains and it is only adequate to support animal life from the same circumstance.

But as it has been before observed oxygen gas is not fit of itself for respira-

-tion and for this purpose is the negative principle nitrogen gas added to weaken the powerfull effects of the oxygen gas or to dilute it as it were. –

The atmosphere (from αηρ, air and σφαιρα a globe) or the Common air as it is emphatically called is well known to be that diaphanous inodorous fluid which surrounds our globe. It is inferred to extend to the distance of 40 or 45 miles in height at the level of the Ocean it is capable to sustain a column of water 35 feet high or one of mercury of the height of 30 inches and it presses with the weight of about 15 ~~inches~~ pounds on every square inch of surface. As we ascend the atmosphere decreases in density in geometrical proportion to equal ascents.

The weight of the atmosphere was ascertained with great care by Mr. Brande from a number of experiments at the Royal Institution he found that 100 cubic inches of air at a mean temperature and pressure of the ~~thermometer~~ and barometer to weigh only 30.199 grs. The atmosphere acts a very important part in equallizing temperature; the great body of air is always in constant motion; over the torrid zones it is expanded by the concentrated rays of the sun; it being expanded its specific gravity would be lessened and consequently it would rise to the superior regions and flow over towards the poles; while the air from the temperate and frigid zones would press forward to fill up

the vacuum. This process still going on and the heated portions coming in contact with colder portions and with various parts of the earths surface which may be hotter or colder is supposed to occasion the various currents of air which we meet with on the earths surface. In this manner also are the different currents at the surface and at various depths of the ocean accounted for; the water of the other parts of the globe mixing with the tropical water. Thus we find a beautifull provision of nature for supporting a uniform temperature. The atmosphere then possesses all those properties which are eminently suited for the comfort and convenience of creation; a deprivation of this fluid would annihilate all nature; all the vegetable and animal creation would cease to exist; the sun would cease to send her cheerfull and invigorating rays on earth; the moon would be blotted from the sight; all nature would pass into darkness and chaos would regain its pristine domain.

The air is accused as the cause of numerous diseases; and it really is so. Sudden cold checking the perspiration will apparently produce almost every form of pyrexia. Partial cold will produce Rheumatism, Pneumonia &c; Damp air, Catarrh. The continued heat of summer occasion bilious disorders and

the cold of winter a return of more active inflammation.

The pure air we breath is often charged with bodies foreign to our comfort and health. We allude to the infinite variety of adventitious bodies which are found floating or mixed as it were with the atmosphere. Every reading man knows whether high or low wet or dry the atmosphere is always in connection with various substances the principle of which are three and they are the only ones that we shall notice in this place.
1. Carbonic acid. 2. Watery Vapour.
3. Principle of Infection; which last has occupied the attention of men for a long time and we shall endeavour to give it some share of our attention.

Carbonic acid, is at all times present in the atmosphere and is estimated by Mr. Dalton not to exceed one thousandth of its bulk. By Van Helmont it was called, gas sylvestre from being produced in vast quantities from burning Charcoal; from its apparent acid properties, aërial acid; and Carbonic acid; and fixed air as readily losing its elasticity and fixing itself in many bodies. It is an invisible and permanently elastic fluid; superior in gravity to the common air and most other aërial fluids. It consists of twenty eight parts of

carbon and seventy-two of oxygen and undoubtedly with some caloric. It is unfit for respiration; easily absorbed by water exceedingly destructive to animal life and produced in great quantities naturally from combustible bodies and many chemical processes. It is found at the bottom of pits and caverns; near Naples there exists the famous Grotto del Cane which is constantly filled with it; it arises from fermenting liquors. It is heavier than the atmosphere 100 cubic inches according to Sir H. Davy at 60° Farenheit weigh 47.11 grains. This gas is powerfully antiseptic destroying the putrefactive decomposition. When introduced into the stomach it is agreeable and grateful and is administered with advantage in some disorders: but though it may be introduced into the stomach and intestines with good effect if breathed into the lungs it is mortal. By the respirations of animals comparatively a large quantity of carbonic acid is formed: this results from the union of the oxygen of the atmosphere with the carbone of the venous blood.

Plants absorb carbonic acid gas and restore in their turn pure oxygen which combining with azote may imperceptibly to our senses renovate the atmosphere. Thus nature very completely restores the various changes in the com-

-stitution of our atmosphere which the dif-
-ferent processes constantly going on may
in her regular course have occasioned.

Watery Vapour. That watery vapour exists
in the atmosphere sometimes requires not the
mind of a philosopher to prove. Every
man of reason knows very well that the
atmosphere very often contains water in
a state of vapour — Now it has been found
that the atmosphere contains watery va-
-pour at all times, at all seasons even in
the coldest weather. It is well known that
beside water; Camphor, Volatile Salt &c if
placed exposed to the air will evaporate; &
even if we view ice exposed to the air still
a portion of it is lost by evaporation and
we are told that the vapour produced in
this way is different from that produced
by boiling namely that it is not so elas-
-tic. This sort of evaporation is called spon-
-taneous in order to distinguish it from that
produced by more obvious sources of heat.

It is the opinion of Mr. Dalton that wa-
-ter existing in the atmosphere constitutes a
distinct and independent atmosphere of its-
-elf: this opinion of Dalton's has been con-
-sidered by many and perhaps with truth
as very erroneous; for as Dr. Henry says it
is certainly more reasonable to suppose that
water whenever it exists as an elastic fluid
neither distinct from or mixed with others, is

maintained as such by me under the same cause viz. the Caloric which enters into it: and not by Chemical solution in any gas or mixture of gases. It is stated that the reason vapour is lighter is that its specific gravity is less than that of water. The explanation which is now generally received is that of Dr. Halley's. He compares the case to a solution of Salt in water considering the air as the water and the vapour as the Salt he still farther states in corroboration of his hypothesis that the collateral circumstances which favour solution are favourable to spontaneous evaporation; such as an increase of temperature, extent of surface, agitation, &c. After the philosophic world had received this explanation of Dr. Halley's a circumstance was remarked which threw over the whole. It was found by putting a cup of water on the plate of an air pump and exhausting all the air from the bell that evaporation would go on and to an extent greater in a given time than when placed in the open atmosphere the evaporation went on under the bell of the air pump untill it was filled with vapour which put a stop to the farther evaporation of the fluid. This simple experiment then proves the inconsistency of Dr. Halley's supposition for in this case the collateral causes which he alludes to as assisting the evaporation in the open atmosphere could not have operated here and another thing

to be attended to was that in the experiment with the air pump the pressure of the atmosphere was taken off the surface of the water.

It is stated that vapour itself exists in the atmosphere in the form of small particles which are vesicular or little hollow globes.

Different portions of vapour are formed at different altitudes and in different latitudes of the earth; in the torrid zones it is most abundant; in the temperate less and in the frigid zones still less vapour is found in the atmosphere; the quantity is however governed very much by the season of the year &c.

In the torrid zones very little rain falls vegetation is supplied by the profuse dew which is formed in the night after the sun passes the horizon of these regions the same quantity of vapour is no longer kept suspended in atmosphere it consequently falls unobserved through the atmosphere to the ground in the form of dew and supplies the place of profuse rains which fall only in the temperate zones. In the northern part of the temperate and in the frigid zones the power of the sun not being strong enough to suspend the water in the form of vapour condenses it which descends to the surface of the earth through the atmosphere in the form of hail, frost, snow &c these are the products of varying nature produced by climate and change of season.

Principle of Infection. It was an object of great interest among the medical world in those days when the constitution of the atmosphere was first discovered to suppose that the cause of some peculiar diseases was owing to a too less quantity of oxygen in certain portions of the atmosphere; it was supposed that in high situations there was more oxygen than in low situations and especially in low marshy places there was a considerable less quantity of it. Now it was very natural for men to draw this inference yet like the fate of all hypotheses it has to fall when opposed by facts. The Chemists turned their attention to this point and to prove wether the atmosphere suffers any deterioration in this respect they collected the air of marshy places; they took it from jails, from prison ships; and from every infectious place and examined it and yet they found the very same principles in the very same quantity always they could detect no less of either the one or the other of the principles composing the atmosphere. A very celebrated Chemical philosopher Gay Lussac ascended to a very great height in a balloon and he took with him a glass jar filled with water; when he had arrived at a very great distance from the earths surface he emptied it and immediately corked

it right; he brought it down with him to his labor-
-atory; he first examined the air of his labor-
-atory that there might be no mistake and
found that it contained the usual proportions of the two ingredients; viz; oxygen and Nitrogen gases. He then examined the air which was contained in the glass jar and he found it to contain the usual quantities to a fraction of a the gases which was in the air of his laboratory.

This then was proof enough to any honest minded man that the cause of an infectious disease was not to be attributed to a too great quantity of oxygen in the atmosphere in any given place.

But when this hypothetical notion had been put down by Chemical investigations; some who wished to explain the subject in a new way gave rent to their imagination in bringing forth the ideal agency of the existence of Inflammable-air; Hydrogen; in the atmosphere as the cause of the pestilential disease Yellow fever We not ~~fine~~ stop to refute this opinion for it has been shown by Davy that hydrogen cannot exist alone for any length of time in the atmosphere without combining with it; and if this be the case and in all probability it is the case; it would soon be diffused equally throughout the whole atmosphere and thereby so diluted as to be

altogether innocuous.

It was the opinion of some that the existence of the nitrous acid vapour in the atmosphere was also the cause of the fatal and pestilential disease Yellow fever and it was furthermore supposed that the exhilerating gas of Davy had some connection is the cause of a pestilential disease; Yet from the experiments of Beddoes and Davy it cannot be supposed to exert any deleterious influence; on the contrary it has the effect of a cordial or invigorating remedy much less a depressing or debilitating effect.—

A very ingenious physician of the South Dr. Sheut advances the doctrine that the infectious disease Yellow fever is dependent on the existence of an aerial fluid in the atmosphere in consequence of a derangement of the Electrical Equilibrium in the air. That when by the occasion of lightening the Electrical Equilibrium is restored it is impossible for the gaseous poison or Yellow fever infection to exist in such a degree as to produce the epidemic Yellow fever. In support of his doctrine the author adduces a number of facts as derived from the medical history and annual meteorological observations for South Carolina. Like many

other disputed points in medical science authors for the most part speak doubtfully as to the precise nature of the infection which produces fevers of various degrees of malignancy: we allude now principally to that fever which has received the name of Yellow fever. – It is now settled by the majority of the learned of the medical world that the greatest quantity of this peculiar principle which has been denominated Marsh miasma; arises from marshy grounds more or less impregnated with putrid exhalations from the decomposition of vegetable substances.

Its activity & producing fevers is promoted by all those circumstances as increase the putrefactive decomposition and promote the exhalation of it; such as heat and moisture. When marshy grounds are at particular seasons of the year covered with water this prevents the miasma from rising in any considerable quantity and of consequence the diseases are fewer and less violent. But when from the great evaporation occasioned by a hot and dry summer; the ground is uncovered leaving it in a swampy state then aquatic plants, trees, &c; undergoing the vegetable decomposition, produce an exhalation which is highly

acilterious.

Although authors appear to agree that marsh miasma is the most general cause of the epidemic yellow fever and give the most undeniable proofs of its effects; yet we are not to conclude marshes to be the only source of this miasma. For every place where there is vegetable matter combined with moisture and heat decomposition of the vegetable matter will necessarily take place and give rise to miasma which may be as prolific in producing its peculiar disease as that arising from a marsh.

From the preceding short view of the nature and origin of miasma it will appear that with the concurrence or absence of particular circumstances tending to increase or diminish its noxious power it must in different times and places exist with various degrees of strength.

Sometimes it is scarcely capable without the conjunction of other causes of producing fever; whilst at other times it may be of so noxious a nature as to affect the most robust in a short time after it has been received; and we can frequently observe intermediate degrees of its malignancy both with regard to the power of the noxious effluvia and also to the force

of the system, from habit, to resist their effects.

The manner in which miasma effects our system so as to prove the cause of disease is not clearly ascertained although physicians of the present day believe that it is taken in by the saliva swallowed and then taken into the stomach or either by Respiration. Some are inclined to one opinion and some to the other. The first opinion that the infectious effluvia mixes with the saliva and afterwards gets into the stomach has been embraced by several physicians of great character and learning; and as a proof of this opinion they find by not swallowing their spittle when attending the diseased they are exempt from the disease;

This appears very plausible; but we never could in our mind be led to suppose that the first impression of the cause operated in this way; for in the very few instances that might be alledged in proof of this hypothesis it may be refferred to peculiarity of constitution &c and not to that of the miasma mixing with the saliva and afterwards getting into the stomach and there fermenting with the contents of the prima via and so causing an increased secretion of bile &c!

That of respiration is more plausible the miasma is taken into the lungs during the act of inspiration it is consequently

brought into contact with the sentient extremities of the nerves and there primarily produce its peculiar impression as a cause producing the disease. By this view of the subject we think the phenomena of the disease can be more rationally accounted for.

It would require Volumes to illustrate and examine the various points in dispute among medical men relative to this peculiar principle of infection as the cause of Yellow fever we have said thus much on this part of the subject the rest we leave to those who have more time knowledge and experience;

Vale.

---

437

435

437

# On the Hydrocele of the Tunica Vaginalis Testis,

an Inaugural Dissertation,

submitted to

The Faculty of Physic

of the

University of Maryland,

by

John Berry,
of Nashville Tenessee,

for the

Degree of Doctor of Medecine,

March 1828.

The term Hydrocele applies to any watery tumour, but the subject of this essay is confined to that disease which is denominated the Hydrocele of the Tunica Vaginalis Testis.

This is a disease from which no period of life is exempt. Not only adults but young children are frequently affected with it, and infants are sometimes born with a congenital Hydrocele. The earliest account of this species of Hydrocele was drawn up by Mr. Viguerie a surgeon at Toulous and communicated to the Academy of Surgeons at Paris. From numerous observations it is said to be much more frequent in Germany, than in England. It is likewise remarked that this species of hydrocele, is frequent and of long continuance in Jewish children.

The parts concerned in the Hydrocele of the Vaginal tunic, are the following. The common bag in which both the testis are enclosed, called the scrotum, which consists of epidermis, skin, and that loose cellular membrane called the dartos, to which may be added the expanded fibre of the cremaster muscle.

The proper coats of the testicle are the tunica

albugenia, and tunica Vaginalis. The former of these immediately invests the vascular portion of the testis, and is that coat which covers them while within the cavity of the ~~tentum~~ abdomen, before they descend into the scrotum. The latter is spread ~~over the opposite side of~~ on the said cavity and is a process of the peritoneum placed ready for the reception of the testicle, when it shall have passed down to its destined situation.

Between the vascular structure of the testicle and the tunica-albugenia there is no cavity, but the external surface of the gland is in every part adherent to, and connected ~~with~~ with, the internal one of the investing coat. The tunica Vaginalis forms a hollow cavity or bag which is unconnected at the superior anterior and lateral parts of the testicle, being at the same time firmly united to its posteriorly ~~part~~, in such a manner, that if the cavity of the tunica vaginalis was to be distended with air, such air would occupy or fill the loose and unconnected part, and produce a tumefaction not unlike hydrocele, while the testis would be found firmly attached to the hinder part of the distended cavity.

Mr. Pott supposed that whatever tends to increase the secretion of fluid in to the cavity of that membrane beyond the due and necessary quantity, or to prevent its being taken up and carried off by the absorbent vessels, contributes to the production of this disease. Murysh entertained the opinion that hydrocele might arise from a varicose state of the spermatic veins. Mr. Hamsden conjectured, that it arose sometimes from an irritated state of the urethra. This disease has been known to originate during an attack of rheumatism, Exposure to cold and the pressure of ill constructed trusses on the spermatic cord, have sometimes been alleged as the cause of this desease. It is also said to be of more frequent occurence among old men and persons who ride much on horse back.

Dropsy in general and this disease often arises from an increased secretion of the arteries, or a diminished action of the absorbents; but dropsical swellings are

generally the result of an increased secretion from the arteries. The proofs are found in the increased vascularity of the membranous surface from which this secretion is poured out, as is seen upon inspection of the parts post-mortem, and also in the change which is found to be produced in membranes of similar structure after long continued dropsies; and in the quickness with which Hydrocele succeeds inflammation of the testis and tunica Vaginalis, Hydrocele is most commonly the result of a relaxation of the arteries from whose patulous mouths an inordinate effusion of fluid takes place. It more frequently arises from this cause than from inflammation. In Hydrocele the absorbent vessels of the spermatic cord, on the diseased side, are said to be very much larger than on the opposite and healthy side. It is sometimes though rarely consequent upon inflammation of the testicle, which subsiding leaves the tunica Vaginalis filled with water. It is generally a local disease but is sometimes connected with an universal hydropic diathesis

## Symptoms and progress.

In general on its first appearance, the tumour is rather round, but as it increases it frequently assumes a pyriforme figure, with its largest extremety downwards; sometimes it is hard, and almost incompressible, somuch so, that in some few instances, it has been mistaken for a induration of the testicle; at other times it is soft and lax, so that both the testicle and fluid surrounding it are easily discoverable. It is perfectly indolent in itself, though its weight sometimes produces a small degree of uneasiness in the back.

Mr Pott states the transparency of the tumour to be the most uncertain sign belonging to it; he says it is a circumstance which does not depend on the quantity colour or consistence of the fluid constituting the disease, somuch as on the uncertain thickness of the containing bag, and the common membrane of the scrotum.

According to Sir A Cooper, upon accurate examination of the swelling, it is found to be transparent. His directs is to examine the tumour in the following way. The room is first to be darkened, the patient must then hold a candle burning brightly close to the side of the scrotum; the surgeon must grasp the posterior part of the swelling so as to render its fore part as tense as possible and then look at the swelling from the side opposite to the candle; placing his feet

hand on the fore part of the scrotum he will immediately discover the transparency. The strong light of the sun falling on the part answers equally well in shewing its transparency.

The quality and consistence of the contained fluid is various It is thin, aqueous, ropy, viscid, limpid, yellow, greenish, brown, bloody, clear, or turbid. It has a distinct fluctuation which may be observed by placing the hand upon one side of the tumour and gently striking it upon the opposite side. When excessively distended it feels hard. The quantity of fluid varies in different cases; a hydrocele of ordinary size contains about eight or ten ounces, but instances are on record in which four and even six pints of water have been drawn off.

The progress of the disease is generally so slow that six or seven months often elapses before the tumour approaches the abdominal ring. This disease is more inconvenient on account of its size and weight, than painful or dangerous. When large and not supported in a bag truss it drags the spermatic cord and creates pain in the loins. It also interferes with every kind of labour in which the lower extremities are concerned, and prevents the individual from riding on horse-back. Among other characters of this case the corrugation of the scrotum disappears by the distention of the parts, and the manner in which the swelling draws the integuments from the penis prevents the full erection of that organ; a circumstance which

is frequently a source of great anxiety to the patient, who is apt to suppose his virility irrecoverably impaired. When the swelling is very large, the penis is completely retracted, and the urine dribbling over the front of the scrotum, is liable to bring on a good deal of inflammation and troublesome ulceration of the parts.

In the Hydrocele of children the testis occupy a lower situation than the same organ in the hydrocele of adult persons, and the swelling appears farther up towards the abdominal ring; however the testis is generally placed two thirds of the swelling downwards, and at the posterior part of the scrotum. Pressure at that part gives the sensation of squeezing the testis. It is a very moveable swelling and does not distend the part much in the course of the spermatic cord; it bends easily upon the abdomen and moves readily in all directions.

## Diagnosis.

Hydrocele may be distinguished from Hernia by the occasional return of the herniab swelling into the abdomen, by the dilation of the hernia in coughing, and by hernia descending from the abdomen. In sarcocele in which the testis is truly diseased, the tumour is more heavy and flatter on the side than hydrocele, and more solid. Much pain is also produced by

squeezing the testis, the epidydymis is often capable of being felt as a distinct tumour; the cord may be traced with facility. There is also great vascularity of the scrotum, and the general health is often impaired. From Hæmatocele we may distinguish it by the greater weight of the former by its want of transparency by its obscure fluctuation, and particularly by its being usually the result of a blow upon the part.

From Cirsocele it may be distinguished by placing the patient in the recumbent posture in which situation by pressure on the distended veins the tumour may be made to disappear. From hydrocele of the cord by the latter disease extending some times above the ring; it is also globular when grasped and raised; it appears of a light blue colour is very transparent and extremely free to the feel unattended with pain; it is merely an inconvenience to the patient from the impression it produces on his mind. Hydrocele may distinguished from Hernia humoralis by the Hernia involving the whole scrotum, by its being attended with much inflammation and generally resulting from gonorrhœa.

## General Treatment.

In general when a hydrocele is left to itself in an adult subject, no cure can be effected by nature alone. In young-

persons the cure is more easy, in children under six years of age nature without any assistance sometimes disperses the complaint. Sometimes the tumour bursts or is ruptured by external violence, or from an accidental effort of the patient, Accidents of this kind sometimes terminate in a radical cure.

Internal remedies have no effect in this disease, unlesp employed in a very early stage, in cases of young children cold affusions of water have been found useful, When this remedy is used it is recommended to be poured on the part out of a tea-pot four or five times a day. Active purging has in some instances effected a speedy cure of this disease however it is rarely found to be useful.

## Palliative Treatment.

Hydrocele is not a painful disease. In many cases, the patient generally prefer submitting to the inconvenience it occasions for some time, rather than have recourse to any surgical operation, However its weight and size finally become so disagreeable that he is obliged to wear a bag-trup, a painful extension of the spermatic cord takes place

and the patients very much annoyed by a troublesome excoriation caused by the friction of the tumour against the sides of the thighs. Hence the greater number of patients are very anxious for relief.

All the anterior superior and lateral parts of the vaginal coat are loose and detached from the albugenia, but on its posterior part those two tunics make one, and consequently the testicle is as it were affixed to the posterior part of the cavity of the sack in hydrocele. This being the state of the parts the operation ought always to be performed in that part of the tumour where the two coats are at the greatest distance from each other and where the fluid must be accumulated in the largest quantity.

The two instruments used for this operation are the common lancet and the trocar. For the purpose of discharging the fluid, the latter instrument is generally preferred, as the canula facilitates the escape of the water and prevents its diffusing itself in the cellular texture of the scrotum, when the hydrocele is small or the

459

An

Inaugural Dissertation

On Tetanus

Submited to an examination

the Faculty of the University

of Maryland

for the degree of Doctor of Medicine

by Joseph N. Stuart

of Virginia

1828.

Tetanus as defined by all authors is a contraction of several of the muscles of greater or less violence and extent with tension and rigidity of the parts affected. Tetanus may be divided according to its severity into acute and chronic. The first is very dangerous, and generally mortal. Chronic tetanus is less dangerous and by reason of its more gradual progress and affords time for the employment of more numerous remedies.

Tetanus may from certain causes occur in every climate with which we are acquainted but it occurs most frequently in the warmest climates and generally in the warmest seasons of those climates. It affects all ages and complexions. The causes from whence it commonly proceeds are cold and moisture applied to the body while it is very warm, and especially the sudden vicissitudes of heat and cold, or by punctures lacerations or other ~~lesion~~ lesions of nerves in any part of the body. Possibly there are many other causes of this dreadful disease; but they are neither distinctly known, nor accurately ascertained.

If the disease proceed from cold it most frequently comes on in a few days after its application; but if it arise from a puncture of a nerve the disease does not commonly come on for many days after the injury has happened, very often when there is neither pain or uneasiness in the part, and frequently when the wound has cicatrized. Sometimes this disease comes on suddenly to a very violent degree, but it approaches more generally very gradually to its most violent state.

In this case it comes on with a sense of stiffness in the back part of the neck, which gradually increas-

ng renders the motion of the head difficult and
tremely painful. As the rigidity of the neck comes on
nd increases there is commonly at the same time a sense
f uneasiness felt about the root of the tongue, which by
degrees becomes a difficulty of swallowing, and at length
an entire interruption of it; when this pain arises all
he muscles of the neck and particularly those on the
back part of it are immediately affected with spasm,
bulling the head very strongly backwards. At the
same time the temporal and masseter muscles which
pon the first approach of the disease were affect-
ed with some rigidity are now affected with violent
pasm, and set the teeth so closely together that
ey admit not of the smallest opening. This is what
commonly has been called Locked-jaw, and often
is the principal part of the disease.
hen the disease has advanced thus far a pain at
the bottom of the sternum returns, and leaves
he patient very frequently; and with it the spasms
f the muscles of the back part of the neck and lower
aw are renewed with great violence, and severe pain.
s the disease proceeds a greater number of muscles
become affected with spasm. After those of the
eck, those along the whole spinal column are
ffected, bending the trunk of the body back-
wards; and this constitutes what has been called
opisthotonos. Both the extensor and flexor muscles
f the inferior extremities are commonly at the

same time affected, and keep the limbs rigidly extended; though the extensors of the head and neck are usually more strongly affected, yet the flexors & those muscles that pull down the lower jaw are often strongly affected with spasm at the same time. During the whole of this disease the abdominal muscles are violently affected with spasm, so that the belly is much retracted and feels very hard. When the disease has made some progress the flexors of the head and trunk are equally affected with the extensors, so as to keep the head and trunk straight and strongly extended, incapable of being moved in any way, to this state the term tetanus has been strictly applied. At this time the arms which were little affected before become rigidly extended; and the whole of the muscles belonging to them being affected with spasm, except those that move the fingers which often in the most violent stage of the disease retain some mobility. The tongue also retains its mobility for a long time; but at length it also becomes affected with spasms which attacking only some of its muscles, forces it sometimes between the teeth, by which it is much injured. At the height of the disease every organ of voluntary motion seems to be affected and among the rest the muscles of the face.

The forehead is wrinkled, the eyes are sometimes

distorted and immoveable in their sockets, the cheeks are drawn towards the ears, & the countenance is expressive of most violent agony. Under these universal spasms the patient does not long remain, for a convulsion soon terminates his existence. In this disease the spasms are every where attended with great pain. The utmost violence of spasm is however not constant, but after subsisting for a short time, the muscles admit of some relaxation, although not so much as can allow the action of their antagonists. This remission of contraction gives a short remission of pain, but neither is of long duration. These violent contractions & pains are removed from time to time, & often times without any evident exciting cause. Every attempt to speak, to move or to swallow, gives occasion to the renewal of the spasms over the whole body. The attacks of the disease are not always but generally attended with febrile action, at the commencement of the disease, the pulse is full & frequent, tense & strong. When the spasms are general and violent, the pulse is contracted, hurried and

irregular; and the respiration is affected in a like manner but during the remission both the pulse and respiration usually return to their natural standard.

Frequently the face is pale, with a cold sweat upon it, and very often the extremities are cold with a cold sweat over the whole body. In this disease the head is seldom affected with delirium, or even confusion of thought, till the last stage - so that the patient is sensible of all the changes of this distressing disease, his pain is unsupportable, he sees the desponding countenances of his attendants and feels as if his departure could not be arrested. In this disease the natural functions are not immediately affected. Vomiting sometimes appears early but it is not continued, the appetite commonly remains good nearly through the whole course of the disease, & the food taken seems to be well digested. The excretions are generally deranged. It is not certain whether costiveness is the effect of the disease or of the opium which is administered, probably the latter as it is used in large quantities. The urine is sometimes suppressed, or it is voided with difficulty and pain. This disease has commonly proved fatal as we are told by Dr Cullen

and fatality might might be considered as a natural consequence as long as the disease was badly treated but since a more proper method has been known and practised, the disease has not struck the patient or the physician with such terror as formerly, and many now are cured. Judging of the tendency of the disease in particular cases, we may remark, that when it arises from a laceration of the nerves it is commonly more violent, than when it is the effect of cold; that the disease which comes on suddenly and advances quickly to a violent degree is always more dangerous, than that which is slower in its progress. This disease often proves fatal before the fifth day, and if the patient has passed this period, he may be supposed to be in greater safety and generally the longer it continues, the safer the disease and consequently the less do we fear the result. However it is to be remarked that many days after the fifth the disease continues to be dangerous; and when it is considerably abated, it is ready to recur again with its former danger. No evacuation or eruption has been known suddenly to check its career, and when it terminates favourably it always recedes gradually, and often a long time elapses before all its symptoms disappear entirely. When the disease is known to arise from the laceration

of a nerve in any part of the body, the first and most important step to be taken towards the cure, is by every means in our power to cut off that part from all communication with the sensorium either by cutting through the nerves in their course, or by destroying the afflicted parts to some distance. We have several instances of great benefit from early amputation of a wounded extremity when tetanic symptoms first appear. This however should not be done unless the injury from which the disease results be inflicted on some of the smaller members such as the fingers or toes; to amputate the thigh would be putting the patient to useless torture. Opium as taught by experience has often ~~produced a~~ proved a very effectual remedy when we attempt to cure the disease by medicine. The remedies have been divided into external and internal, when the disease arises from the injury of a nerve it will be necessary to cut down and divide it, or to apply stimulating remedies or warm emolient poultices, & at the same time to give Laudanum, under this treatment the disease will be frequently removed. Both the warm & cold bath have been strongly advocated, but the cannot be relied on alone. Caustic has been often used successfully in traumatic tetanus. Of the internal remedies opium is the best & should

never be forgotten in the treatment of tetanus and cannot be limited. it should be given as soon as you suspect that the disease is approaching. wine or brandy may be give in very large quantities, in this disease it is almost impossible to produce intoxication; The whole class of antispasmodics have be used the oil of Amber is among the best & if used in larger quantity would more often prove successful it should be given in the dose of half a dram. When opium is administered it should be given in small & frequently repeated doses, at the intervals of 2 or 3 hours as the violence of the symptoms may require, it seldom induces ~~torpor or sleep~~ stupor or sleep which it does so often does under other circumstances when much smaller doses have been taken. therefore we should not be sparing in its use.

Purgatives have been administered & recommended by very respectable authority in this case - Dr Hamilton of Edinburg represents them as entitled to our confidence. But if they be useful it must be in cases of tetanus resulting from internal irritation, where the disease originates in the alimentary canal. In such cases probably emetics combined with purgatives might be useful, for besides assisting in the removal of irritation they would contribute by their nauseating quality to the relaxation of spasm. Blood letting has been practised with advantage in this disease in plethoric subjects, regard must be paid to the state of the system in this as in all other complaints. Blisters were formerly in this disease, but it is asserted that they were

constantly hurtful & are now never employed.

Mercury in its turn was tried, & report says with success; & is now one of the principal remedies in use. This, with opium & wine, appear to be the only remedies, upon which we can confidently rely, in the treatment of tetanus. Of these, the two latter retard the progress of the disease, preventing exhaustion, & death, from violence of the spasm, until the former has had time to eradicate it entirely, by converting the tetanic into the mercurial action. Opium & wine being palliative rather than curative remedies are not sufficient of themselves without mercury while without their aid mercury is too slow in its operation to arrest the rapid march of the disease. The usual & best preventatives of tetanus from external injuries consist in the conversion of punctured into incised wounds by means of the scalpel & the excitement of suppuration in lacerated wounds.

These things if done immediately after the injuries have been received & before the occurrence of tetanic ~~symptoms~~ spasms are of great importance but they are worse than trifling if delayed untill the disease shall have made its appearance. When the tetanic diathesis is fairly formed all local applications to the injured parts are unavailing even the amputation of a limb has been found insufficient to arrest its progress. Should circumstances exist to prevent the dilation of a punctured wound the application of caustic alkali & sinapisms to the part are the best substitutes. Whatever is calculated to excite common inflammation is useful at this time. For the production of suppurative action in lacerated wounds spirits of turpentine, or a solution of the muriate of mercury followed by poultices frequently repeated are the best applications. When from a neglect of preventative remedies, or in despite of their use tetanus has made its appearance, a different course of practise must be commenced and pursued diligently. To the state of the

system a strict and steady attention must be paid. If the patient be plethoric and the excitement run high, or if the latter, without the former blood must be drawn and the operation repeated as often as circumstances may require; if costiveness prevail & any irritation be suspected in the bowels purgatives are to be administered as in other diseases; but as has been mentioned above as soon as the condition of the system will admit them, wine opium, & mercury are to be given according to their effects on the system without regard to quantity. They should be given until the system has forcibly felt their action or they will be unavailing. Opium should be administered in injection in doses of from five to ten grains & repeated at short intervals until the spasms are overcome and some soporific effect produced; applying in the mean time to various parts of the body quantities of mercurial ointment, when the power of swallowing has for a long time ceased, nutritive clysters should be substituted. After surgical operations, as a prophylactic salivation, & the use of the tincture of opium mixed with the dressings has been employed to prevent symptomatic tetanus with success.

481

983

985

An
Inaugural Dissertation
on
Sclerus
Submitted to the examination
of
the provost
the Trustees and Medical Professors
of the
University of Maryland
For the degree of Doctor of Medicine
by
Edward McCeney of
Anne Arundel County Maryland
March 27th 1828

To

Samuel Baker M. D.

Professor of Materia Medica

in the

University of Maryland

Permit me sir to dedicate this dissertation to you as a small tribute of gratitude and esteem for the polite attention and benefits derived from your instructions during the prosecution of my medical studies under your guidance they were always gratefully received and will ever be remembered by your former pupil Edward McCeney

# Essay on Icterus



Before entering upon the subject of Icterus I propose to make a few physiological remarks on the liver Considering the size of this organ in all animals that possess it and how generally it is possessed being common to all red blooded animals there can be no doubt that it is of great importance in the animal economy. Whether it serves to separate any excrementitious material from the blood or produce any other effect upon it has not yet been decided Certainly its most obvious office is the secretion of the bile which in most animals in its passage from the liver to its place of destination the duodenum communicates by means of a duct with a pear shaped reservoir denominated the vesicula fellis or gall bladder in which it is suffered to accumulate And as ~~the~~ it ~~symptoms~~ is generally believed that the symptoms of Jaundice depend upon an obstruction to the free egress of the bile from the excretory ducts of the liver into the alvine canal and its return into the circulation Although such is the case in general yet there have been instances in which there were a copious absorption and

return of bile into the blood and at the same time a redundant flow of it into the alimentary canal However the most general causes of Icterus may be referred to mechanical obstructions to the biliary ducts Many disputes have arisen concerning the manner in which the bile is resorbed into the system but it is now generally believed that it is taken up by the absorbents of the gall bladder and ducts Hence it is that jaundice may be produced by any thing obstructing the passage of the bile into the duodenum Biliary concretions appear to be the most frequent cause of this disease With respect to their formation we are yet unacquainted The liver appears in many instances to be sound but in other cases it is undoubtedly diseased and from the morbid secretion it may become more disposed to chrystalize From the analysis of biliary concretions they appear to contain most of the ingredients of the bile From their peculiar chrystalized structure we are led to believe that something more is necessary to their formation than the mere inspissation of the bile but what that consists in physiologists have not yet been able to detect Sudolence of body & anxiety of mind appear

particularly to predispose to their formation therefore they are much more frequently found in women in the upper walks of life and men much addicted to sedentary occupations especially after the more active period of life Haller noticed the frequent occurance of biliary concretions among criminals whose deaths had been preceded by long confinement but the anxiety of mind also might contributed to their formation as long as biliary concretions remain undisturbed in the gall bladder for they are generally formed there they appear to produce little or no inconvenience but from violent compression or from irritation the gall bladder and ducts contract and thus the offending substance will gradually be moved in the direction where least resistance is offered this will of course be toward the mouth of the duct not solely because the duct gradually enlarges but because there is constantly an impulse given from the accumulated secretion from behind. When biliary concretions of large size are thus thrust into the ducts they produce the most excruciating pain to which the human fabric is liable and it is generally refered to the pit of the stomach

corresponding to the opening of the duct into the duodenum and shooting from thence towards the spine accompanied at intervals with rigors and nausea During the paroxysm the respiration is hurried with great anxiety restlessness and occasionally with delirium followed by extreme languor and faintness with vomitting & occasionally with spasmodic ~~tw~~ twitchings in different parts of the body Profuse sweats often break out apparently from the severity of the pain In the intervals of the paroxysms there is generally a fullness and soreness in the epigastric and right hypochondriac regions The patient generally experiences most relief when his body is bent forward The paroxysm will continue to recur at intervals until the concretion makes its escape into the alvine canal which it will do in some instances within a few minutes in others it will be for weeks in passing along the duct thus obstructing the passage and producing the disease which in some instances will continue for a considerable time after the cause has ceased to act Biliary concretions of such large size have sometimes passed as to make physicians doubt whether they did not make their way by ulceration into some part of the

intestinal canal which is highly probable as there are cases where they have been known to make their way through the walls of the abdomen

Preternatural viscidity of the bile has often been adduced as a frequent cause of jaundice It appears that this state of the bile may take place from a variety of causes From the rumora of the bile in the gall bladder and ducts produced by particular positions of the body from pregnancy and from vascular turgescence of the liver itself and at the same time the absorbents of the gall bladder and ducts still continuing to act if not increased in action will take up the more alienate particles of the bile and thus render it viscid Augmented secretion from the gall bladder and ducts has also been assigned as a frequent cause and also a too viscid consistence of the bile in its secretion in the liver This species of jaundice is most frequently met with in autumn and it generally commences very insidiously There is in this variety a diminution of appetite restlessness disturbed sleep and a disinclination to exertion of any kind together with the general symptoms of Icterus

Inflammation appears to be a very frequent cause of Jaundice by preventing the free discharge of the bile probably by a thickening of the parietes of the ducts thus lessening their calibre It may take place from a variety of causes it may either extend from the intestines or from the liver itself from cold and especially from taking very large potations of cold water when the body is much heated The symptoms attending this species of obstruction are those attending general inflammation tenderness of the epigastrium and generally commences with a chill It has been questioned by some whether Jaundice is ever produced by mere spasm of the ducts themselves. The arguments in favour of this opinion are that it has been a frequent attendant on Hysteria and other spasmodic diseases and that the attack is sometimes very transitory and from dissections there has been discovered no mechanical obstructions whatsoever in the ducts and also without any appearance of biliary concretions in the faeces It occurs more frequently in persons of very irritable habits from sudden emotions of the mind The passions have a remarkable effect on the secre-

-tion of the bile and they may also cause a spasm of the ducts themselves The symptoms attending spasmodic constrictions of the ducts are sometimes as violent as when bilian concretions are impacted in the ducts The spasm indeed appears to be the cause of the pain in both cases. Enlargements of the neighbouring organs is not an unfrequent cause of this disease such as schirrhus of the pancreas scrophulous glands tuberculi of the liver may ob-strup the parietes of the ducts as to prevent the discharge of the bile and thus produce the disease In this variety the pain is not so acute and does not come on suddenly general emaciation attended with soreness and hardness in the epigastric region Jaundice is also described to have been produced by mechanical distention of the duodenum by closing the valve of the duct Jaundice arising from this cause appears to be peculiar to infants and is very easy of removal It only requires the exhibition of some mild aperient medicine to clear the bowels of the impacted viscid meconium

Having spoken of the principal causes which

give rise to the disease I shall now enumerate some of the most characteristic symptoms Jaundice generally commences with a want of appetite a sense of lassitude and languor depression of spirits disinclination to exertion of any kind Indigestion and general emaciation In some cases there is a great propensity to sleep in others wakefulness The dejections are pale or clay coloured with the absence of the feculent smell As the disease increases the yellow colour makes its appearance showing itself first in the tunica albuginea The urine becomes high coloured with a yellowish sediment so as to be capable of imparting the same colour to linen The yellow colour appears not to be confined to the skin alone but is diffused over the whole body so that the bones have been found tinged yellow for a very long time after the disappearance of the disease There is generally great itching over the whole body The bile appears to be diffused in the whole mass of fluids so that the secretions are affected even the blood itself is said to possess a yellow colour.

as jaundice may arise from so many different causes
some of which it is impossible to discover during the
patients life and some which may be considered
incurable we ought therefore to ascertain if possi-
-ble the nae nature of the cause which has given
rise to the disease which can in general be done
by attending to the various circumstances which
present themselves for certainly such a discrimi-
-nation must be of the greatest importance when
it is considered that the mode of treatment must
be varied according to the nature of the cause by
which the disease is produced Whether the passage
of the bile be obstructed by biliary concretions or
spasmodic constrictions the plan of treatment
to be adopted must be nearly the same In the
passage of biliary concretions of very large size
through the common duct they very frequently
produce from their great distention not oly very
acute pain but also when long continued a consi-
derable degree of inflammation to guard against which
it will often be necessary in full plethoric ha-
bits to take away blood proportionable to the

state of the pulse and severity of the pain This should be succeeded by the use of the warm bath warm fomentations together with opiates The opiate should be given in the solid form for the stomach will often be so irritable as to reject liquids of every kind Two or three grains of opium should be given at first and if relief is not obtained in half an hour the same quantity should be repeated with the intention of facilitating the passage of biliary concretions or viscid bile emetics have been used with great advantage During the act of vomitting the diaphragm and abdominal muscles concurrantly contract and the whole of the abdominal viscera are thus forcibly pressed upon which must necessarily cause the gall bladder and ducts to pour out their contents freely For this purpose the antimonial emetics are preferable because they are less apt to be rejected and also excite a much stronger stimulus After the operation of emetics the bowels should be freely opened by some purgative medicine Exercise has also been useful particular

-ly that of riding on horse back by the concussion which it gives to the liver frictions and the like have been used with the same view Purgatives have been found very useful in some cases of Jaundice especially in those cases where the disease arises from a too viscid state of the bile in consequence of a torpid condition of the liver In this case mercury must be used to change the condition of the Liver In cases of infants where the passage is obstructed by viscid meconium purgative medicines must be principally relied upon The nitro-muriatic acid bath hath been much recommended in cases of Icterus It should consist of three ounces of diluted aqua regia to every gallon of water the strength of the bath should never be greater at any time for otherwise it may produce a very troublesome rash and give a yellow hue to whatever part is exposed to its action Dr Scott used it with decided advantage in almost all cases dependant on a morbid secretion of bile there has been a great variety of other remedies recommended in Icterus such as the alkalies electricity the seed of the common hemp &c but none of them much used at the present time

591

515

# An,

Inaugural, Essay

On the,

Causes of Sudden death,

And the means of preventing it,

Submitted to the examination,

Of the,

Medical Proffessors,

Of the,

University of Maryland,

For the degree of,

Doctor of Medicine,

On, the day of, 1828,

By Daca, Wilson,

Of Alabama,

# On the,
# Causes of Sudden death,
# And the,
# Means of preventing it,

517

Death is the inevitable lot of all mankind but there are few persons whose business or whose conduct in life, do not render it a greater evil to themselves and their families when it takes place suddenly than when it occurs after a previous indisposition. To point out the principal causes of the sudden and unexpected extinction of life, and the means of preventing it shall be the subjects of the following dissertation. It will be necessary that I promise to addopt the opinion of the professor of the Institutes that life consists in such a peculiar organization of matter, as to be capable of producing by certain impressions, those motions and sensations which have been hypothetically ascribed, by some, unites to the mind and by others to an independent vital principle. Death of course consists in the distruction of that peculiar organization of matter, or an incapacity

with those impressions which produce the motions and sensations of life. The causes which produce this incapacity in the matter of the human body, and thereby induce sudden death, act more or less certainly, according as they are more or less combined, according to the nature of the parts of the body on which they act, according as the system is in a state of healthy excitement, or more or less predisposed to death, by previous debility, or by a previous disease. They appear to induce death by a sudden destruction of the excitement of the system from the excessive force of impression or the sudden abstraction of them, whereby the system is rendered incapable of performing its natural offices.

I shall briefly enumerate the causes, which, by inducing the above changes in the system, bring on sudden death. Extreme heat and extreme cold. Both these act with most certainty upon old and sickly people. Dr. Husham remarks, that old people are often found dead in their beds, after an extreme cold or warm night; and their are few physicians who have not had patients suddenly and unexpectedly snatched out of their hand by the extremes of heat and cold. Fatal effects have likewise followed or rather been observed when heat and cold in their extremes, have followed each other

in a rapid succession. The rays of the sun acting upon the head have often produced death, by means of what is called by the French coup de soleil:" and cold, combined with moisture, has so often produced it in the form of Apoplexy, as in some instances to create a belief that disease had assumed the character of an epidemic. Sudden changes in the weight of the atmosphere, some cases of kind are mentioned by writers. Certain matters contained in the air. These are several of the gases particularly azote or according to modern nomenclature nitrogen gas, the carbonic Acid, and inflamable Air. The black at Calcutta will be a lasting monument of the deadly effects of azote or respired Air. Many people perish from sleeping in unventilated cabins and rooms, in consequence of Carbonic Acid eliminated from burning coals, and it is well known that miners are often destroyed by breathing the inflamable Air which is discharged from subterranious caverns. To the noxious airs some have added certain vegetable poisons which are said to float in the atmosphere: but since the rejection of the fable of the East India Upas tree, sudden death from this cause has been discarded from Medicine. The Air in the vicinity of the Rhus Radicans has been known to induce [illegible] and to inflame the skin, but I believe in no instance

has it suddenly extinguished the vital. To this head belongs the ~~vital~~ Miasmas which exist in the atmosphere from the putrefaction of Animal and vegetable substances. The history of the plague in the Old world, and of the yellow fever in the United States, furnish many proofs of sudden death from this cause. This is not the place to inquire whether those diseases or death can be induced by a matter secreted in a morbid body, and conveyed into a healthy one, sudden death has sometimes been brought on by entering a small and filthy room, crowded with persons ill with the plague! But the matter which induces it is not secreted. It is like the original Miasms, the offspring of putrefaction, and in these cases, derived from stagnated excretions from the bodies of sick people. As I am only relating facts at present, I hope I shall be excused from deciding on the long agitated question of the part of the body, on which the Airs and Miasms, that have been mentioned, primarily and chiefly act, in producing their deadly effects. Some of them certainly act upon the body, through the medium of the lungs; some of them upon the whole nervous system through the medium of the brain; And some of them upon the sanguiferous system through the medium of the stomach. It is possible, Miasms may convey death to the body through the avenues of all the parts that have

been mentioned. I have said nothing of sudden death from contagions. They are extremely limited in their number, and seldom so powerful, in their first impressions as to extinguish life. Cases of sudden death from the Small pox, Measles and the hooping cough, are I believe rarely to be met with in the records of medicine. Lightning is often a cause of sudden death. It seems to act primarily upon the nervous system, but the blood and all the muscles of the body appear to partake of its forcible impression. Hence the quick putrefaction of the body after death from this cause. Sudden and violent emotions of the mind. Anger has in instances produced death terror has often produced it. Soldiers has often been found among the slain in battle whose bodies have been found free from wounds. Their deaths have been ascribed to the wind of cannon balls; Is it not probable they were occasioned by the violent impression of terror upon their systems: Trifling wounds it is well known, have now and then produced death, from being combined with a sudden paroxysm of fear. The passions of fear and grief have occasionly induced sudden death, but it is more common for them to produce it by a previous chronic disease. Under this head we may bring in as a cause of

sudden death, a sudden sense of gill and shame. Of
this their are several instances to be met with in
practical authors. Great bodily exertion in labours lifting
weights and in marching have sometimes suddenly dest
royed life. Fatigue or the reduction of excitement of the
system in marching more certainly induces sudden death,
if it be combined with excessive heat. The histories of the
the French and English campaigns in Egypt furnish a
number of instances of Soldiers expiring on the public
road, from the combined effects of the rays of the
sun, and great fatigue acting upon their bodies. Great
bodily pain has sometimes induced sudden death. The col
o often kills by excessive pain. Mr Hunter relates an insta
ce of the pain occasioned by the extirpation of a diseas
which producing death. In fevers sudden death is sometimes
brought on by the inability of the bloodvessels to react
in the chilly fit; by convulsions and by the extreme de
bility which follows. The termination of a paroxysm
of pron. affections of the head, from Gout and the
frequents causes of sudden death. The most common
of these are presence of water or blood upon the brain
inducing apoplexy. Death is sometimes the effect of a
sudden effusion of blood or water in the lungs, constituting
what has been called by Dr Rush in his lectures the apoplexy
of the lungs. A Spasm of the heart, produced by the gout,
as in several instances, induced the sudden extinction
of life. Certain poisons and many substances of an in
digestible nature, also cold liquors, when the body is
preternaturally heated, received into the stomach by their
sympathy with the head, heart or bloodvessels, induce
sudden death. It has sometimes followed a Spasm
in the stomach from gout, also a rupture of that im
portant viscus. The discharge of pus from the liver into
the thorax, and into the abdomen, has often hurried persons

suddenly out of life. Worms in the alimentary canal have now and then suddenly produced death, especially in children, by exciting apoplexy or general convulsions. The sudden rupture of an aneurism, in the large blood vessels near the heart, has often become an instantaneous extinct of human life. It has passed away with nearly equal rapidity from a rupture of the bladder and intestines. It has been induced by syncopes from partial and general causes. The sudden reduction of the excitement of the system, by Hemorages, by parturition, by a large stool, by a copious discharge of matter from an abscess and perhaps, by the infliction of pain, has in sundry instances brought on sudden death. Derangement producing suicide. I have ascribed this cause of sudden death to a cause, for such is its contrariety to the natural principle of action in man, that I believe it rarely takes place in the perfect exercise of his reason. I shall say nothing of the causes of sudden death from drown external causes wounds and accidents as being foreign to the subject, and briefly to mention the means of preventing it. I shall confine my self only to such causes as are within the power of medicine.

Art. 3rd To prevent death from the storms of heat; cool drinks and a situation should be advised. As it frequently occurs in the night, and during sleep, in warm weather, by recommending to old and weakly persons to constantly on a mattress, to prevent it from cold; warm but light bed cloths, should be prescribed; and as death makes its first ~~approach~~ approach through the feet, great pains should be taken to keep them warm, by means of warm socks, wrapped in warm flannels, or stone jugs filled with warm water. The fatal effects of the stroke of the sun are to be prevented by constant and gentle exercise, when persons are exposed to its rays, and by protecting the head by

an umbrella, or a deep crowned hat, covered with white linnen or paper. The Egyptians, says Mr. Duvar, use a turban for this purpose. It is true it accumulates the heat of the head but this heat he remarks, is less, and far more tolerable, than the heat of the sun, which in the climate of Egypt, is often between 129° and 139° degrees. Within these degrees of heat says Dr. Rush, I have seen the natives of Bengal rowing their boats, on the river Hoogly, exposed to a meridian sun and bare headed, without experiencing any ill effects. The only precaution they used, was to keep their heads constantly wet by pouring on them large gourds of water. By these means the neck, shoulders, and chest which with the head were exposed to the sun, were kept cool from the evaporation of the water. During the prevalence of the winds and that kind of weather which disposes to sudden death from apoplexy, great care should be taken to lessen the predisposition to it, by temperate diet and gently opening physic. To prevent the fatal effects of the gases, that have been mentioned, places which are supposed to contain them should not be entered, until they have been examined by a lighted candle. The extinction of its light, is a sign of its containing fixed or carbonic gas. A flame or explosion excited by the flame of the candle, indicates the presence of inflamable air. The deadly effects of the miasms from putrefied vegetable and animal substances, are to be prevented by avoiding the places which generate them, by lessening the predisposition to be affected by them, by means of abstinence gently opening physic, and a temporary issue, excited by a blister on one of the legs, or limbs. Of the efficacy of these preventives there are many proofs in the history of pestilential fevers and diseases. The means of preventing or avoiding sudden death from lightning, belong to naturals not to medical philosophers. The preventions of death from violent emotions

f the mind, should be sought for in those books which inc
culcate the subjection of all its faculties, and operations to the
dictates of reason, and religion. Labour and exertion disp
portioned to the strength of the body, should be careful
y avoided by all who wish to avoid or escape a sudden
-ap out of the World. The combined influence
heat and fatigue upon life from labour and ma
chinery may be obviated by counter stimuli. Ardent spirits
ave generally been resorted to for this purpose, but garlic
raw Onions are to be preferred to them. By the constant
e of these cordial vegetables, the hebrew nation were enabled
sustain the labour of making bricks, in the open Air
posed to the intense heat of the climate of egypt.
he deadly effects of heat and fatigue were prevented
r a part of the British army in the same country,
uring the late war, by allowing the Soldiers, but a small
uantity of water. Similar advantages are derived from
bstaining from water altogether, by the Indians of North
America, during the day, in their long and fatiguing
marches, in hot weather. Death from extreme pain may
e prevented by bleeding. Opium, hunger and giving vent to
its sensations by cries and groans. Of the efficacy of
hunger in easing pain, Dr. Priestly relates the following: In
avelling through germany, he passed through a village, where a
riminal was exposed to torture once a week, in order to extor
om him a confession of his accomplices in his crimes, three days
fore he suffered he avoided tasting any food. The sensation
hunger thus induced was so painful, and at the same time
harmless in its effects upon life, as to be over balanced
pain and danger of the tortures inflicted upon him. unex
ted death in the chilly fit of a fever, should be avoided
(ann drinks; from excessive force of reaction in the blood

helps by bleeding; from convulsions, by remedies suited to the stage
the disease in which they occur, and from the debility which
llows a paroxysm of fever by cordial medicines, drinks and
iments. Sudden death in persons predisposed to it from apop
y should be obviated by temperance moderate exercise,
en bowels; not suppressing usual evacuations, nor changing usual
its; by occasional bleedings when the premonitory signs of
tigo; and head ach take place; by avoiding tight ligatures
ound the neck, by sleeping with the head a little elev
d, and upon a mattress instead of a feather bed
d by never sleeping upon the back, but alternately
on each side Dr Rush mentions several cases of palsy,
does Dr Potter in his lectures, a disease produced by a
ore feeble operation of the causes which produce Apoplexy
Rush mentions that patients sleep upon their left side.
sudden extinction of life by the Apoplexy of the lungs
Dr Rush would call it and considers appropriate and
itimate, by Dr Potter in his lectures, by poisonous and
digestible matters received into the stomach, by worms
the alimentary canal; and by spasm of the heart,
uld be prevented by the usual remedies for that purpose,
tioned in the book of Medicine.

den death from cold liquors, taken into the stomach when the
y is preternaturally heated, has been often prevented
plunging or rather conducting off the heat of the body
plunging the hand and feet into cold water, or by wash
the mouth with it, or by grasping the vessel contain
the liquors provided it be made of a metal or
y kind of earth. 12th Persons predisposed to
ut; Spasms in the stomach or heart, should
void the remote or exciting cause of Gout; but
hen attacked by it, should flee to Laudanum, untill
advice of a physician can be obtained.

When danger is apprehended of the sudden discharge of pus from the liver into the thorax or abdomen, it should if possible, be diverted externally, by means of a caustic. Pills which emulge the bile from the liver into the stomach and bowels, would perhaps have the same effect, for its purulent contents, the ducts which lead to the duodenum would ~~perhaps~~ probably yield to the agitation excited by a vomit, sooner than any other part of the liver. 14th Sudden death from aneurisms has often been prevented by a temperate diet, tranquility of body and mind, and by small and frequent bleedings. When sudden death is apprehended from syncope, great pains should be taken to avoid the remote and exciting causes and to strengthen the system by tonic remedies. The sudden extinction of life from sudden depletion, or the sudden collapse from distention, or cessation of pain, should be obviated by vicarious stimuli of other kinds: when they cannot be prevented by other means. 15th When there is reason to apprehend sudden death from suicide, persons suspected of it should be narrowly watched, and all the means of death should be removed from their vicinitude should be prevented, and cheerful society or cheerful glass, should be advised. The excitement of strong emotion of terror and pity has prevented it in numerous instances. 17th In cases of sudden death from any of the causes which have been mentioned, it is probable that resuscitation might be effected by the usual means, when judiciously applied, In the use of them

shall only deliver the following directions: 1st begin
with such as are of a gentle nature, and gradually
resort to such as are more powerful, in all cases where
the cessation of the functions of life has been induced
by causes which leave the excitability of the system
in an accumulated state. 2nd In cases where by great
exertions to preserve life, the excitability of the system
has been expended, let no stimulus be applied for some
time after the cessation of apparent life. By observing
this delay, time will be given for the accumulation
of some of the feeble remains of excitability in the sys
tem. 3rd In addition to the common remedies employed
to effect resuscitation, feeble or loud sounds applied
to the ear, are calculated to produce the most sal
utary effects. Life often lingers longest on the
ears, so says Dr Rush in his lectures on asphyxia.
It has been proved by the shrieks of distressed persons
sometimes calling their apparently dead friends to
life. By availing ourselves of this fact, a new
and powerful remedy, may be added to those which
are now in common use for promoting resuscitation.
That the motions of life are destroyed last in the
brain (and of course in the vicinity of the ears),
I think highly probable from the placid or gloomy
countenance which succeeds death. They are probably
the effects of the pleasant or painful operations of
the mind, which survive the extinction of life in
other parts of the body.

An
Inaugural Dissertation
on
Inflammation
Submitted
To the
Examination of the Provost
and
Professors of the University of Maryland.
By their
Friend
William H. Kirk
of
Lancaster County
Virginia
on the 3rd
of
April
1838

In the prosecution of our medical studies, there is no subject which presents itself to our view more interesting than inflammation, or one for which we as surgeons are more frequently called to treat, therefore a perfect acquaintance becomes necessary.

- Inflammation has been divided by surgeons into acute and chronic, simple, or complicated with disease; acute or healthy inflammation has been often termed phlegmon, and is of one kind only. Whilst malid inflammation is subdivided into many kinds as erysipelatous, scrofulous, venereal &c. &c.

= Phlegmon, or healthy inflammation is that which has for its object the restoration of diseased parts. Malid inflammation is that which has some morbid action added to that of unhealthy, such as in carbuncles &c.

= The general characteristicks of phlegmonous inflammation are an increased sensibility of the part, pain, preternatural redness, accompanied with tumefaction, and throbbing. The increased redness is supposed to arise from and increase flow of blood through the small arteries of the part, sometimes it may be dependant upon a generation of new vessels. Samuel Cooper says that the swelling is dependant upon a dilitation of the vessels, to a plethoric state of the arterial system & veins to the exudation of coagulation of lymph into the

interstices of the cellular substance and also to an interruption of the absorption.
With regard to the increased heat this seems to be imaginable, as the degree of heat scarcely ever surpasses the natural temperature of the body, but we may readily account for this when we remember the extreme sensibility of the nervous system, by which they convey false impressions.
The remote causes of inflammation are numerous consisting in those of a mechanical or chemical nature such as bruises, wounds, splinters, acrid substances &c. Inflammation sometime arises as consequent on some febrile disease acting upon the system and when this is the case is called symptomatic sometimes it seems to arise without any obvious cause & then it is commonly called spontaneous but improperly, as there can be no effect produced without there be a cause to produce that effect.
The proximate cause of inflammation has been a source of much theoretical speculation, it was supposed by the ancients that it was dependant on some morbid condition of the fluids, Whilst the moderns as Cullen and others, that it is dependant on a change of the vessels & not a disease of the fluids, The absurdity of the antients may be clearly seen from a local inflammation, for were it dependant upon a disease of the fluids, we could have no inflamma-

tion however slight without bringing the whole system into a state of disease. It seems more probable to depend upon a disease of the vessels, from the fact that new vessels are sometimes formed, this may be proved by tumours, which may be extirpated without occasioning much hemorage when at the same time were we to cut into the liver, the hemorage would be great. It seems to me that all inflammation must be dependant upon an increased action of the vessels in the part affected, for it cannot be doubted that there is more blood sent to the part when inflamed than when in a state of health, this seems clear from the circumstance, that if we cut into an inflamed part it will bleed much more freely, than when in a natural state, neither can this be done by any increased action of the heart, for the heart sends its blood equally to the whole system, but from this circumstance it will seem, that besides the elasticity peculiar to the arteries, themselves they have also a power of dilating and contracting of themselves, we can scarcely explain on any other principals, the increased flow of blood through the inflamed part.

It has been thought by some that it is dependant upon a spasm of the extreme vessels of the part, but I think that an effusion of lymph besides the increased size of the vessels would be sufficient to overturn an such an idea as this.

A very peculiar circumstance attending phlegmonous inflammation is that let it be situated, where it may or in what part soever it will always point externally. If inflammation invades the sockets of the teeth the matter will point externally, immediately in the vicinity of the cheek, than on the inner side of the alveolar process.

Inflammation differs in there varieties in there progress and there termination, as they differ in there situation, position, structure, and functions of parts. Parts through which the blood circulates freely can undergo inflammation much better than are situated in opposite condition, except when the part is essential to life, and then though the part may be possessed with a great deal of vascularity still it is owing to the uniformity which is required for the support of health dependant upon a sound condition. Having now given the symptoms and causes on what we think the most reasonable, from some consideration and considerable study, we will now endeavour to give the treatment as recommended by the best modern surgeons. In the treatment of inflammation the first to be considered is the removal of the remote cause for if there be a splinter or any foreign substance lodged in the flesh we should first extract that, and then we should have an eye to the proximate cause, it will seem at once from what has been said that

our principal motive must be to remove that cause this is to be effected by those remedies which is necessary to subdue the increased action of the enlarged vessels.

## Bleeding

As we have advanced the idea that inflammation is dependant upon an increased action of the blood vessels in the part affected; it will not be doubted that blood letting forms one of the most prominent agents in the reduction of this action, this should be done either by topical or general bleeding. Topical bleeding seems in many cases to be more efficacious than general, when the part affected is local and does not bring the system into concern, topical bleeding is more benificial than general, but when any part is affected which is essential to life then we should use general and topical both, thus where the brain is affected & the inflammation is not speedily subdued death will be the inevitable consequence, we should use general blood letting when the part is not possessed with that power of restoration, which is peculiar to the mucous, as in tendons about the joints, also when suppuration would be the result and that perhaps to the destruction of the patient. Opthalmia is a case of this kind although it would not destroy the patients life yet the organ itself would be destroyed, General blood letting should not admitted when the inflammation

is trivial and excited in an old debilitated subject. When it takes place in a young and plethoric we should use the remedies as early in the disease as possible, when the action runs very high we should always use general bleeding first, to make our topical remedies act the more effectual, we should draw the blood from a large surface, as to the quantity of blood to be taken we should be governed entirely by circumstances, the time of the disease when taken and the general excitement in the system, which may always be told by the strongness and fulness of the pulse.

Purging

These form the next grand feature in the catalogue of our remedies for the cure of inflammation, they should be of the saline kind, they not only assist blood letting in diminishing the circulation, but they act by producing a secretion they may be used even when blood letting cannot be admitted, those most commonly used is the Sul. of Soda, Sul. of magnesia, and the nitrate of potash.

Antimonials.

Nauseating doses of antimonials preparations are sometimes very beneficial they act by removing the offensive dryness of the skin, and from the sympathy which exist through the nervous system diminishing diminishing the action of the inflamed vessels this is accomplished

by the nervia which is produced occasioning a tempo
rary debility and through these the progress of the in
flammation is subdued —

## Opium

When there is great constitutional irritation together
with great pain opium may be employed, although
in the most cases of inflammation opium cannot
be administered in account of its stimulating effe
cts.

## Diet

This in the cure of inflammation is very essential. the
patient should be kept from all animal food, from
fermented & Spirituous liquors, a neglect on the part
of the Surgeon, to enforce, and the patient to adhere
to a strick antiphlogistic regimen, often is the cause
of the inflammation terminating in that way which
brings reproach upon our science and in some
cases to the destruction of our patients

## Cold applications

This remedy has been recommended from the know
ledge that heat or rather apparent heat has allways
been the concomitants of inflammation, and by
the abstraction of that heat the part effected will be
restored, this is done by keeping up a constant evap
oration on the surface. The remidies which are
generally employed by Surgeons are of an astringent
or sedative a nature, such as the acetate of Lead dissolv

ce in water. Mr Cooper recommends the following formula acetate of lead ℥ss vinegar ℥iv. distilled water ℔ii, but I am inclined to think that the water & vinegar contain ace the virtues, because I cannot perceive what the virtues of the lead can consist as it cannot have any specific action on the part, but the whole benefit must be owing to the evaporation which takes place, as the modus operandi of the lead has not as yet been explained. Doctor Smith, the professor of surgery in this institution has recommended an evaporating ~~poultice~~ lotion composed of spirit and water both equal parts, which has much the advantage of the lead & water in subduing the inflammation, as I have had the opportunity of seeing the wonderful effects during the last winter in a case in our Infirmary of a man who had punctured his foot with a nail, when he was brought into the Infirmary, there was a violent inflammation taken place and in the course of forty eight hours all the symptoms were almost subdued. Again I have seen the formula which Cooper recommends fail entirely

Warm application

This has been also recommended but it seems strange that any two remedies so very opposite to each other should be employed in the same case with advantage but so it is experience has shown that it is so, and it is thought by some surgeons that it is dependent

on some peculiar idiosyncracy of the patients constitution, or some difference in the construction in the part effected. Of the warm application emollient poultices have been the most frequently employed & the best is that made of Linseed meal and warm water this is made by putting the meal into the water untill they are of a proper consistance to form a poultice and then add a little oil to prevent the poultice from becoming hard

## Termination of Inflammation

Inflammation after continuing for sometime, disappears without any apparent cause, or exciting a disposition to form pus, or runs on to such a high state of inflammation as to terminate at loss in the entire death of the part. The first is called resolution, the second called suppurating, & the third and last is called mortification.

## First or Resolution

The most fortunate way in which inflammation can terminate is resolution this consist in a spontaneous cessation of all the symptoms, a subsiding of the redness pain swelling, throbbing & heat of the part

## Second or Suppuration

Suppuration is the next most frequent termination of inflammation, this consist in the formation of pus which modifys the symptoms rather than an entire cessation of them

## Mortification

The worst manner in which inflammation can terminate is in mortification but luckily it is the most rare; this consists in the entire death of the part. I shall conclude with the subject here leaving a particular description of those three varieties alone leaving that they be long more strictly to Symptomatic fever than to this place.

563

An

Inaugural Dissertation

on

Necrosis

presented to the Faculty of the

University of Maryland

for

the degree of Doctor of Medicine

by

George F. Carmichael

of

Virginia

1828.

# A Dissertation on Necrosis.



The strict interpretation of this term is the death of some portion of the bony structure, but is applied by surgeons to a disease characterized by peculiar symptoms, a frequent termination of which is the death or exfoliation of a portion of bone.

This disease commences with an acute inflammation of the bone itself or of its periosteum, accompanied with violent pain which is not at first referred to the immediate seat of the attack but to the nearest joint, it however locates itself finally on the part inflamed. The symptomatic fever which generally occurs at the same time with the pain, is of a highly inflammatory character. The termination of the local affection, is most frequently in suppura-

tion. The matter is deposited between the external periosteum and the bone. If the disease be located in the shafts of the long bones, at the same time, that the matter forms beneath the external periosteum, there is a corresponding collection between the internal lining and the bone. This circumstance has been repeatedly ascertained by the use of the trephine.

Shortly after the attack the limb becomes swollen which is uniform until the matter escapes from beneath the periosteum, when the swelling is more circumscribed and confined to the immediate seat of inflammation. At this period there is a subsidence of the severe pain & fever in some degree, but they do not leave the patient entirely. The death of the bone in this disease

is not attributable to any peculiar malignity of the inflammation, but to the vessels which give nourishment to it, being destroyed by the intervention of matter between it & its periosteum.

When the matter contained beneath the periosteum, has made its way to the surface and that within the cavity of the bone, has escaped through fissures in the same, relieving the parts from irritation and pressure the symptomatic fever in a great degree subsides, if however the collection be considerable and the distruction of bone extensive, hectic fever supervenes, showing a continuance of irritation tho' changed, as to its character.

The process instituted for the purpose of removing the dead bone, depends upon the

extent of injury, for instance if merely a small portion, situated on the shaft of a long bone be necrosed, granulations will shoot from the sound and living bone, and as occurs in the mortification of the soft parts separate the dead from the living portion, and finally urge it through the opening previously formed, and disengage it from the body, this is more likely to happen when the soft parts have been freely divided at the beginning of the disease.

When the whole circumference is destroyed it is covered by an osseous structure which envelops the sequestra and is attached to the sound & living portions not however embracing it closely, and thus giving the limb an enlarged appearance,

Necrosis is almost exclusively confined to young persons seldom or never occurring under five or over twenty two.

No bone in the human system is exempt from this disease, long bones however are more liable, to its attacks, and of these the tibia more particularly, the articulations are never assailed by this kind of inflammation in the first instance, sometimes however the joints are secondarily affected from the disease being communicated to them along the shafts of the long bones, where it first existed.

The line of seperation is often accurately defined by the junction of the shaft & epiphysis of the bone, the articular portion remaining uninjured, it sometimes happens however that the disease reaches the joint and matter is formed when it becomes necessary to amputate the limb to

and the life of the patient,

Necrosis is not always confined to one bone, it may occur simultaneously in remote parts or successively which is more common. In regard to general prognosis a very great majority of patients survive the attack after long confinement and protracted suffering, it sometimes prove fatal but generally early in the attack from the high degree of symptomatic fever and constitutional irritation.

Diagnostic Symptoms according to Dr Smith. ~~The pain~~ "As the disease say this Gentleman is an acute inflammation characterized by the peculiar vital properties of the parts affected in any of the symptoms must be analagous to those of other inflammatory affections, he has often known it to be mistaken and for a considerable time treated for acute Rheumatism, even

altho' suppuration may have been observed, It frequently happens that when the disease fix-es on one of the long bones and near its ex-tremity the pain is complained of in the ad-jacent joint, it is not long however confined to the joint, but fixes itself in the inflamed part, This circumstance of pain referred to the joint has often caused the disease to be denominated Rheumatism."

The pain experienced in Necrosis is extreme-ly acute, unremitting, and not much in-fluenced by the motions nor position of the limb, The pain is often antecedent to the swelling, and when the latter first occurs it is generally diffused over a considerable part of the limb, especially below the part affected, The surface is rather firm to the touch but the skin is not discoloured till after matter is formed and advanced tow-

ards the surface.

The symptomatic fever is coeval with the pain both usually occurring on the same day, the pulse both frequent and quick the stroke sudden and acting small and hard to the touch, At first the patient has occasional chills but when he complains of a sensation of cold the skin to another person feels hot. The pain is so violent that he gets little or no sleep, during the night he is often delirious, The tongue is furred with a soft white coat, the face is not flushed but rather pale with the exception of occasional red spots on the cheeks, The appetite for food is lost the thirst considerable the stomach and bowels are not so much affected as in other febrile diseases.

It has been before observed that the violence of the pain in the early stage of the disease

has caused Necrosis to be sometimes confounded with Rheumatism. Most, even of its early symptoms however are very different from those of that disease, the symptomatic fever and constitutional irritation come on sooner after the local attack and are much more severe, the pulse smaller, harder, and much more easily compressed, and finally suppuration occurring removes all ambiguity. Necrosis moreover usually attacks at that period of life, when rheumatism is not liable to occur. It is distinguishable from Typhus by the local affection, by the pulse which is harder and less easily compressed and by its not being attended with so much stupor.

Causes. The inflammation which produces necrosis is often excited by blows and injuries inflicted upon the limbs, sudden suppression of

perspiration, and emersion in cool water when the body is warm, it however often attacks without any obvious exciting cause, by which the diathesis is developed or concentrated upon a particular part.

Treatment. In the treatment of this disease the first step, as in all other inflammatory diseases should be to effect a resolution, The remedies however, which are most generally employed as venesection, cathartics emetics blisters, evaporating lotions, cataplasms &c, have rarely succeeded in preventing suppuration, the most efficient remedies as directed and practised by Professor Smith, of N. Haven is a free incision of the soft parts, and periosteum down to the bone through the whole extent of the inflammation, thus reducing the disease to a simple incised wound, which go to

be treated accordingly.

In the second stage of the disease when matter has formed beneath the periosteum the cure may still be effected without the loss of bone. In this stage of the disease an incision should be made thro' the soft parts as before directed ~~and~~ the periosteum divided as far as its seperation from the bone, and a portion of the bone be cut out or perforated in the denuded part down to the medullary substance, that the matter collected between that substance and the bone, may escape and thus the death of the bone prevented. The best instrument for perforating the bone is a small trephine.

In the third stage when the matter has made its escape through the periosteum and obtained a lodgment in the soft parts, the treatment is precisely the same

as in the second stage, but the favourable result is by no means so certain, as the circulation in it, may have been too long destroyed, or the seperation of the dead and living bone commenced; nevertheless however as in the former case the incission in the soft parts should be made, taking care, to avoid important parts, such as tendons, nerves, and bloodvessels, this will not only give free vent to the matter contained within, but facilitate the expulsion of the bone, if its distruction should have been extensive,

The general treatment consists in cooling purgatives nauseating doses of Antimony opium sufficient to allay irritation and procure rest.

End